

Rational Associahedra

Drew Armstrong

University of Miami
www.math.miami.edu/~armstrong

Enumerative Combinatorics
MFO, March 2014

Plan

1. Given any $x \in \mathbb{Q}$ define the **Catalan number** $\text{Cat}(x) \in \mathbb{Z}$.
2. Given any $x \in \mathbb{Q}$ with $x > 0$ define the **associahedron** $\text{Ass}(x)$.
3. Given any $x \in \mathbb{Q}$ with $x > 0$ define **parking functions** $\text{PF}(x)$.
4. Have lunch!

Plan

1. Given any $x \in \mathbb{Q}$ define the **Catalan number** $\text{Cat}(x) \in \mathbb{Z}$.
2. Given any $x \in \mathbb{Q}$ with $x > 0$ define the **associahedron** $\text{Ass}(x)$.
3. Given any $x \in \mathbb{Q}$ with $x > 0$ define **parking functions** $\text{PF}(x)$.
4. Have lunch!

Plan

1. Given any $x \in \mathbb{Q}$ define the **Catalan number** $\text{Cat}(x) \in \mathbb{Z}$.
2. Given any $x \in \mathbb{Q}$ with $x > 0$ define the **associahedron** $\text{Ass}(x)$.
3. Given any $x \in \mathbb{Q}$ with $x > 0$ define **parking functions** $\text{PF}(x)$.
4. Have lunch!

Plan

1. Given any $x \in \mathbb{Q}$ define the **Catalan number** $\text{Cat}(x) \in \mathbb{Z}$.
2. Given any $x \in \mathbb{Q}$ with $x > 0$ define the **associahedron** $\text{Ass}(x)$.
3. Given any $x \in \mathbb{Q}$ with $x > 0$ define **parking functions** $\text{PF}(x)$.
4. Have lunch!

What is a Catalan Number?

What is a Catalan Number?

Seemingly Bizarre Convention (It's Not)

Given $x \in \mathbb{Q} \setminus \{-1, -\frac{1}{2}, 0\}$ there exist **unique coprime integers** $a, b \in \mathbb{Z}$ with $0 < a < |b|$ or $0 < b < |a|$ such that

$$x = \frac{a}{b-a}.$$

(Note that $0 < x \iff 0 < a < b$.) We will always identify $x \leftrightarrow (a, b)$.

Examples: Given $1 \leq n \in \mathbb{N}$ we have

What is a Catalan Number?

Seemingly Bizarre Convention (It's Not)

Given $x \in \mathbb{Q} \setminus \{-1, -\frac{1}{2}, 0\}$ there exist **unique coprime integers** $a, b \in \mathbb{Z}$ with $0 < a < |b|$ or $0 < b < |a|$ such that

$$x = \frac{a}{b-a}.$$

(Note that $0 < x \iff 0 < a < b$.) We will always identify $x \leftrightarrow (a, b)$.

Examples: Given $1 \leq n \in \mathbb{N}$ we have

What is a Catalan Number?

Seemingly Bizarre Convention (It's Not)

Given $x \in \mathbb{Q} \setminus \{-1, -\frac{1}{2}, 0\}$ there exist **unique coprime integers** $a, b \in \mathbb{Z}$ with $0 < a < |b|$ or $0 < b < |a|$ such that

$$x = \frac{a}{b-a}.$$

(Note that $0 < x \iff 0 < a < b$.) We will always identify $x \leftrightarrow (a, b)$.

Examples: Given $1 \leq n \in \mathbb{N}$ we have

$$x = n \leftrightarrow (n, n+1)$$

What is a Catalan Number?

Seemingly Bizarre Convention (It's Not)

Given $x \in \mathbb{Q} \setminus \{-1, -\frac{1}{2}, 0\}$ there exist **unique coprime integers** $a, b \in \mathbb{Z}$ with $0 < a < |b|$ or $0 < b < |a|$ such that

$$x = \frac{a}{b-a}.$$

(Note that $0 < x \iff 0 < a < b$.) We will always identify $x \leftrightarrow (a, b)$.

Examples: Given $1 \leq n \in \mathbb{N}$ we have

$$x = \frac{1}{n} \leftrightarrow (1, n+1)$$

What is a Catalan Number?

Seemingly Bizarre Convention (It's Not)

Given $x \in \mathbb{Q} \setminus \{-1, -\frac{1}{2}, 0\}$ there exist **unique coprime integers** $a, b \in \mathbb{Z}$ with $0 < a < |b|$ or $0 < b < |a|$ such that

$$x = \frac{a}{b-a}.$$

(Note that $0 < x \iff 0 < a < b$.) We will always identify $x \leftrightarrow (a, b)$.

Examples: Given $1 \leq n \in \mathbb{N}$ we have

$$x = -n \leftrightarrow (n, n-1) \quad (\text{need } n \geq 2)$$

What is a Catalan Number?

Seemingly Bizarre Convention (It's Not)

Given $x \in \mathbb{Q} \setminus \{-1, -\frac{1}{2}, 0\}$ there exist **unique coprime integers** $a, b \in \mathbb{Z}$ with $0 < a < |b|$ or $0 < b < |a|$ such that

$$x = \frac{a}{b-a}.$$

(Note that $0 < x \iff 0 < a < b$.) We will always identify $x \leftrightarrow (a, b)$.

Examples: Given $1 \leq n \in \mathbb{N}$ we have

$$x = -\frac{1}{n} \leftrightarrow (1, -n+1) \quad (\text{need } n \geq 3)$$

What is a Catalan Number?

Definition

For each $x \in \mathbb{Q} \setminus \{-1, -\frac{1}{2}, 0\}$ we define the **Catalan number**:

$$\text{Cat}(x) = \text{Cat}(a, b) := \frac{1}{a+b} \binom{a+b}{a, b}.$$

Claim: This is an integer. (Proof postponed.)

Example:

$$\text{Cat}\left(\frac{5}{3}\right) = \text{Cat}\left(\frac{5}{8-5}\right) = \text{Cat}(5, 8) = \frac{1}{13} \binom{13}{5, 8} = 99.$$

What is a Catalan Number?

Definition

For each $x \in \mathbb{Q} \setminus \{-1, -\frac{1}{2}, 0\}$ we define the **Catalan number**:

$$\text{Cat}(x) = \text{Cat}(a, b) := \frac{1}{a+b} \binom{a+b}{a, b}.$$

Claim: This is an integer. (Proof postponed.)

Example:

$$\text{Cat}\left(\frac{5}{3}\right) = \text{Cat}\left(\frac{5}{8-5}\right) = \text{Cat}(5, 8) = \frac{1}{13} \binom{13}{5, 8} = 99.$$

What is a Catalan Number?

Definition

For each $x \in \mathbb{Q} \setminus \{-1, -\frac{1}{2}, 0\}$ we define the **Catalan number**:

$$\text{Cat}(x) = \text{Cat}(a, b) := \frac{1}{a+b} \binom{a+b}{a, b}.$$

Claim: This is an integer. (Proof postponed.)

Example:

$$\text{Cat}\left(\frac{5}{3}\right) = \text{Cat}\left(\frac{5}{8-5}\right) = \text{Cat}(5, 8) = \frac{1}{13} \binom{13}{5, 8} = 99.$$

Classical Cases

When $b = 1 \pmod a$ we have ...

- ▶ *Eugène Charles Catalan (1814-1894)*

$(a, b) = (n, n + 1)$ gives the **good old Catalan number**:

$$\text{Cat}(n) = \text{Cat} \left(\frac{n}{(n+1) - n} \right) = \frac{1}{2n+1} \binom{2n+1}{n}.$$

- ▶ *Nicolaus Fuss (1755-1826)*

$(a, b) = (n, kn + 1)$ gives the **Fuss-Catalan number**:

$$\text{Cat} \left(\frac{n}{(kn+1) - n} \right) = \frac{1}{(k+1)n+1} \binom{(k+1)n+1}{n}.$$

Classical Cases

When $b = 1 \pmod a$ we have ...

- ▶ *Eugène Charles Catalan (1814-1894)*

$(a, b) = (n, n + 1)$ gives the **good old Catalan number**:

$$\text{Cat}(n) = \text{Cat} \left(\frac{n}{(n+1) - n} \right) = \frac{1}{2n+1} \binom{2n+1}{n}.$$

- ▶ *Nicolaus Fuss (1755-1826)*

$(a, b) = (n, kn + 1)$ gives the **Fuss-Catalan number**:

$$\text{Cat} \left(\frac{n}{(kn+1) - n} \right) = \frac{1}{(k+1)n+1} \binom{(k+1)n+1}{n}.$$

When $b = 1 \pmod a$ we have ...

- ▶ *Eugène Charles Catalan (1814-1894)*

$(a, b) = (n, n + 1)$ gives the **good old Catalan number**:

$$\text{Cat}(n) = \text{Cat} \left(\frac{n}{(n+1) - n} \right) = \frac{1}{2n+1} \binom{2n+1}{n}.$$

- ▶ *Nicolaus Fuss (1755-1826)*

$(a, b) = (n, kn + 1)$ gives the **Fuss-Catalan number**:

$$\text{Cat} \left(\frac{n}{(kn+1) - n} \right) = \frac{1}{(k+1)n+1} \binom{(k+1)n+1}{n}.$$

Symmetry about $x = -1/2$

Definition

By definition we have $\text{Cat}(a, b) = \text{Cat}(b, a)$, which implies that

$$\text{Cat}(x) = \text{Cat}(a, b) = \text{Cat}(b, a) = \text{Cat}(-x - 1).$$

This implies that for $x \in \mathbb{Q} \setminus \{-1, -\frac{1}{2}, 0\}$ we have

$$\text{Cat}\left(\frac{1}{x-1}\right) = \text{Cat}\left(\frac{x}{1-x}\right).$$

We will call this the **derived Catalan number**:

$$\text{Cat}'(x) := \text{Cat}\left(\frac{1}{x-1}\right) = \text{Cat}\left(\frac{x}{1-x}\right).$$

Symmetry about $x = -1/2$

Definition

By definition we have $\text{Cat}(a, b) = \text{Cat}(b, a)$, which implies that

$$\text{Cat}(x) = \text{Cat}(a, b) = \text{Cat}(b, a) = \text{Cat}(-x - 1).$$

This implies that for $x \in \mathbb{Q} \setminus \{-1, -\frac{1}{2}, 0\}$ we have

$$\text{Cat}\left(\frac{1}{x-1}\right) = \text{Cat}\left(\frac{x}{1-x}\right).$$

We will call this the **derived Catalan number**:

$$\text{Cat}'(x) := \text{Cat}\left(\frac{1}{x-1}\right) = \text{Cat}\left(\frac{x}{1-x}\right).$$

Symmetry about $x = -1/2$

Definition

By definition we have $\text{Cat}(a, b) = \text{Cat}(b, a)$, which implies that

$$\text{Cat}(x) = \text{Cat}(a, b) = \text{Cat}(b, a) = \text{Cat}(-x - 1).$$

This implies that for $x \in \mathbb{Q} \setminus \{-1, -\frac{1}{2}, 0\}$ we have

$$\text{Cat}\left(\frac{1}{x-1}\right) = \text{Cat}\left(\frac{x}{1-x}\right).$$

We will call this the **derived Catalan number**:

$$\text{Cat}'(x) := \text{Cat}\left(\frac{1}{x-1}\right) = \text{Cat}\left(\frac{x}{1-x}\right).$$

Symmetry about $x = -1/2$

Definition

By definition we have $\text{Cat}(a, b) = \text{Cat}(b, a)$, which implies that

$$\text{Cat}(x) = \text{Cat}(a, b) = \text{Cat}(b, a) = \text{Cat}(-x - 1).$$

This implies that for $x \in \mathbb{Q} \setminus \{-1, -\frac{1}{2}, 0\}$ we have

$$\text{Cat}\left(\frac{1}{x-1}\right) = \text{Cat}\left(\frac{x}{1-x}\right).$$

We will call this the **derived Catalan number**:

$$\text{Cat}'(x) := \text{Cat}\left(\frac{1}{x-1}\right) = \text{Cat}\left(\frac{x}{1-x}\right).$$

Symmetry about $x = -1/2$

Definition

Given $0 < x$ (i.e. $0 < a < b$) note that we have

$$\text{Cat}'(1/x) = \text{Cat}\left(\frac{1}{(1/x) - 1}\right) = \text{Cat}\left(\frac{x}{1 - x}\right) = \text{Cat}'(x).$$

We call this **rational duality**:

$$\text{Cat}'(x) = \text{Cat}'(1/x).$$

In terms of coprime $0 < a < b$ this translates to

$$\text{Cat}'(a, b) = \text{Cat}'(b - a, b).$$

This will appear later as **Alexander duality** of rational associahedra.

Symmetry about $x = -1/2$

Definition

Given $0 < x$ (i.e. $0 < a < b$) note that we have

$$\text{Cat}'(1/x) = \text{Cat}\left(\frac{1}{(1/x) - 1}\right) = \text{Cat}\left(\frac{x}{1 - x}\right) = \text{Cat}'(x).$$

We call this **rational duality**:

$$\text{Cat}'(x) = \text{Cat}'(1/x).$$

In terms of coprime $0 < a < b$ this translates to

$$\text{Cat}'(a, b) = \text{Cat}'(b - a, b).$$

This will appear later as **Alexander duality** of rational associahedra.

Symmetry about $x = -1/2$

Definition

Given $0 < x$ (i.e. $0 < a < b$) note that we have

$$\text{Cat}'(1/x) = \text{Cat}\left(\frac{1}{(1/x) - 1}\right) = \text{Cat}\left(\frac{x}{1 - x}\right) = \text{Cat}'(x).$$

We call this **rational duality**:

$$\text{Cat}'(x) = \text{Cat}'(1/x).$$

In terms of coprime $0 < a < b$ this translates to

$$\text{Cat}'(a, b) = \text{Cat}'(b - a, b).$$

This will appear later as **Alexander duality** of rational associahedra.

Symmetry about $x = -1/2$

Definition

Given $0 < x$ (i.e. $0 < a < b$) note that we have

$$\text{Cat}'(1/x) = \text{Cat}\left(\frac{1}{(1/x) - 1}\right) = \text{Cat}\left(\frac{x}{1 - x}\right) = \text{Cat}'(x).$$

We call this **rational duality**:

$$\text{Cat}'(x) = \text{Cat}'(1/x).$$

In terms of coprime $0 < a < b$ this translates to

$$\text{Cat}'(a, b) = \text{Cat}'(b - a, b).$$

This will appear later as **Alexander duality** of rational associahedra.

Euclidean Algorithm

Observation

Given $0 < a < b$ coprime, we observe that

$$\text{Cat}'(a, b) = \frac{1}{b} \binom{b}{a} = \begin{cases} \text{Cat}(a, b-a) & \text{for } a < (b-a) \\ \text{Cat}(b-a, a) & \text{for } (b-a) < a \end{cases}$$

This allows us to define a sequence

$$\text{Cat}(x) \mapsto \text{Cat}'(x) \mapsto \text{Cat}''(x) \mapsto \dots$$

which is a **C**ategorification of the Euclidean algorithm.

Euclidean Algorithm

Observation

Given $0 < a < b$ coprime, we observe that

$$\text{Cat}'(a, b) = \frac{1}{b} \binom{b}{a} = \begin{cases} \text{Cat}(a, b - a) & \text{for } a < (b - a) \\ \text{Cat}(b - a, a) & \text{for } (b - a) < a \end{cases}$$

This allows us to define a sequence

$$\text{Cat}(x) \mapsto \text{Cat}'(x) \mapsto \text{Cat}''(x) \mapsto \dots$$

which is a **C**ategorification of the Euclidean algorithm.

Euclidean Algorithm

Observation

Given $0 < a < b$ coprime, we observe that

$$\text{Cat}'(a, b) = \frac{1}{b} \binom{b}{a} = \begin{cases} \text{Cat}(a, b - a) & \text{for } a < (b - a) \\ \text{Cat}(b - a, a) & \text{for } (b - a) < a \end{cases}$$

This allows us to define a sequence

$$\text{Cat}(x) \mapsto \text{Cat}'(x) \mapsto \text{Cat}''(x) \mapsto \dots$$

which is a **C**ategorification of the Euclidean algorithm.

Euclidean Algorithm

Example: $x = 5/3$ and $(a, b) = (5, 8)$

Subtract the smaller from the larger:

$$\text{Cat}(5, 8) = 99,$$

$$\text{Cat}'(5, 8) = \text{Cat}(3, 5) = 7,$$

$$\text{Cat}''(5, 8) = \text{Cat}'(3, 5) = \text{Cat}(2, 3) = 2,$$

$$\text{Cat}'''(5, 8) = \text{Cat}''(3, 5) = \text{Cat}'(2, 3) = \text{Cat}(1, 2) = 1 \quad (\text{STOP})$$

Euclidean Algorithm

Example: $x = 5/3$ and $(a, b) = (5, 8)$

Subtract the smaller from the larger:

$$\text{Cat}(5, 8) = 99,$$

$$\text{Cat}'(5, 8) = \text{Cat}(3, 5) = 7,$$

$$\text{Cat}''(5, 8) = \text{Cat}'(3, 5) = \text{Cat}(2, 3) = 2,$$

$$\text{Cat}'''(5, 8) = \text{Cat}''(3, 5) = \text{Cat}'(2, 3) = \text{Cat}(1, 2) = 1 \quad \textbf{(STOP)}$$

My Dream / Crazy Idea

My Dream / Crazy Idea

Suggestion

Extend the function $\text{Cat} : \mathbb{Q} \rightarrow \mathbb{N}$ analytically to the upper half plane.



Pause


Well, that was fun.

The Prototype: Rational Dyck Paths

The Prototype: Rational Dyck Paths

- ▶ Consider the “Dyck paths” in an $a \times b$ rectangle.


Example $(a, b) = (5, 8)$



The Prototype: Rational Dyck Paths

- ▶ Again let $0 < x = a/(b - a)$ with $0 < a < b$ coprime.


Example $(a, b) = (5, 8)$



The Prototype: Rational Dyck Paths

- ▶ Let $\mathcal{D}(x) = \mathcal{D}(a, b)$ denote the set of Dyck paths.

Example $(a, b) = (5, 8)$



The Prototype: Rational Dyck Paths

Theorem (Grossman 1950, Bizley 1954)

For a, b coprime, the number of Dyck paths is the Catalan number:

$$|\mathcal{D}(x)| = \text{Cat}(x) = \frac{1}{a+b} \binom{a+b}{a, b}.$$

- ▶ Claimed by Grossman (1950), "Fun with lattice points, part 22".
- ▶ Proved by Bizley (1954), in *Journal of the Institute of Actuaries*.
- ▶ *Proof:* Break $\binom{a+b}{a, b}$ lattice paths into cyclic orbits of size $a+b$. Each orbit contains a unique Dyck path.

The Prototype: Rational Dyck Paths

Theorem (Grossman 1950, Bizley 1954)

For a, b **coprime**, the number of Dyck paths is the Catalan number:

$$|\mathcal{D}(x)| = \text{Cat}(x) = \frac{1}{a+b} \binom{a+b}{a, b}.$$

- ▶ Claimed by Grossman (1950), "*Fun with lattice points, part 22*".
- ▶ Proved by Bizley (1954), in *Journal of the Institute of Actuaries*.
- ▶ *Proof*: Break $\binom{a+b}{a, b}$ lattice paths into cyclic orbits of size $a+b$. Each orbit contains a unique Dyck path.

The Prototype: Rational Dyck Paths

Theorem (Grossman 1950, Bizley 1954)

For a, b **coprime**, the number of Dyck paths is the Catalan number:

$$|\mathcal{D}(x)| = \text{Cat}(x) = \frac{1}{a+b} \binom{a+b}{a, b}.$$

- ▶ Claimed by Grossman (1950), *"Fun with lattice points, part 22"*.
- ▶ Proved by Bizley (1954), in *Journal of the Institute of Actuaries*.
- ▶ *Proof:* Break $\binom{a+b}{a, b}$ lattice paths into cyclic orbits of size $a+b$. Each orbit contains a unique Dyck path.

The Prototype: Rational Dyck Paths

Theorem (Grossman 1950, Bizley 1954)

For a, b **coprime**, the number of Dyck paths is the Catalan number:

$$|\mathcal{D}(x)| = \text{Cat}(x) = \frac{1}{a+b} \binom{a+b}{a, b}.$$

- ▶ Claimed by Grossman (1950), *"Fun with lattice points, part 22"*.
- ▶ Proved by Bizley (1954), in *Journal of the Institute of Actuaries*.
- ▶ *Proof:* Break $\binom{a+b}{a, b}$ lattice paths into cyclic orbits of size $a+b$. Each orbit contains a unique Dyck path.

The Prototype: Rational Dyck Paths

Theorem (Grossman 1950, Bizley 1954)

For a, b **coprime**, the number of Dyck paths is the Catalan number:

$$|\mathcal{D}(x)| = \text{Cat}(x) = \frac{1}{a+b} \binom{a+b}{a, b}.$$

- ▶ Claimed by Grossman (1950), *"Fun with lattice points, part 22"*.
- ▶ Proved by Bizley (1954), in *Journal of the Institute of Actuaries*.
- ▶ **Proof:** Break $\binom{a+b}{a, b}$ lattice paths into cyclic orbits of size $a+b$. Each orbit contains a unique Dyck path.

The Prototype: Rational Dyck Paths

Theorem (Armstrong 2010, Loehr 2010)

- ▶ The number of Dyck paths with k vertical runs equals

$$\text{Nar}(x; k) := \frac{1}{a} \binom{a}{k} \binom{b-1}{k-1}.$$

Call these the **Narayana numbers**.

- ▶ And the number with r_j vertical runs of length j equals

$$\text{Krew}(x; \mathbf{r}) := \frac{1}{b} \binom{b}{r_0, r_1, \dots, r_a} = \frac{(b-1)!}{r_0! r_1! \dots r_a!}.$$

Call these the **Kreweras numbers**.

The Prototype: Rational Dyck Paths

Theorem (Armstrong 2010, Loehr 2010)

- ▶ The number of Dyck paths with k vertical runs equals

$$\text{Nar}(x; k) := \frac{1}{a} \binom{a}{k} \binom{b-1}{k-1}.$$

Call these the **Narayana numbers**.

- ▶ And the number with r_j vertical runs of length j equals

$$\text{Krew}(x; \mathbf{r}) := \frac{1}{b} \binom{b}{r_0, r_1, \dots, r_a} = \frac{(b-1)!}{r_0! r_1! \dots r_a!}.$$

Call these the **Kreweras numbers**.

The Prototype: Rational Dyck Paths

Theorem (Armstrong 2010, Loehr 2010)

- ▶ The number of Dyck paths with k vertical runs equals

$$\text{Nar}(x; k) := \frac{1}{a} \binom{a}{k} \binom{b-1}{k-1}.$$

Call these the **Narayana numbers**.

- ▶ And the number with r_j vertical runs of length j equals

$$\text{Krew}(x; \mathbf{r}) := \frac{1}{b} \binom{b}{r_0, r_1, \dots, r_a} = \frac{(b-1)!}{r_0! r_1! \dots r_a!}.$$

Call these the **Kreweras numbers**.

The Classical Associahedron

Definition

Let $n \geq 0$ and consider a convex $(n+2)$ -gon C . Let $\text{Ass}(n)$ be the abstract simplicial complex with

- ▶ vertices = chords of C
- ▶ faces = noncrossing sets of chords of C
- ▶ maximal faces = triangulations of C

Theorem (Milnor, Haiman, C. Lee, etc.)

$\text{Ass}(n)$ is a polytope.

The Classical Associahedron

Definition

Let $n \geq 0$ and consider a **convex** $(n + 2)$ -gon C . Let $Ass(n)$ be the abstract simplicial complex with

- ▶ vertices = chords of C
- ▶ faces = noncrossing sets of chords of C
- ▶ maximal faces = triangulations of C

Theorem (Milnor, Haiman, C. Lee, etc.)

$Ass(n)$ is a polytope.

The Classical Associahedron

Definition

Let $n \geq 0$ and consider a **convex** $(n + 2)$ -gon C . Let $Ass(n)$ be the abstract simplicial complex with

- ▶ vertices = chords of C
- ▶ faces = noncrossing sets of chords of C
- ▶ maximal faces = triangulations of C

Theorem (Milnor, Haiman, C. Lee, etc.)

$Ass(n)$ is a polytope.

The Classical Associahedron

Definition

Let $n \geq 0$ and consider a **convex** $(n + 2)$ -gon C . Let $Ass(n)$ be the abstract simplicial complex with

- ▶ vertices = chords of C
- ▶ faces = noncrossing sets of chords of C
- ▶ maximal faces = triangulations of C

Theorem (Milnor, Haiman, C. Lee, etc.)

$Ass(n)$ is a polytope.

The Classical Associahedron

Definition

Let $n \geq 0$ and consider a **convex** $(n + 2)$ -gon C . Let $Ass(n)$ be the abstract simplicial complex with

- ▶ vertices = chords of C
- ▶ faces = noncrossing sets of chords of C
- ▶ maximal faces = triangulations of C

Theorem (Milnor, Haiman, C. Lee, etc.)

$Ass(n)$ is a polytope.

The Classical Associahedron

Definition

Let $n \geq 0$ and consider a **convex** $(n + 2)$ -gon C . Let $Ass(n)$ be the abstract simplicial complex with


- ▶ vertices = chords of C
- ▶ faces = noncrossing sets of chords of C
- ▶ maximal faces = triangulations of C

Theorem (Milnor, Haiman, C. Lee, etc.)

$Ass(n)$ is a polytope.

The Classical Associahedron

- ▶ Example: Here is $\text{Ass}(4)$.



The Classical Associahedron

Theorem (Euler, 1751)

The f -vector and h -vector of $\text{Ass}(n)$ are given by the **Kirkman numbers**

$$\text{Kirk}(n; k) = \frac{1}{n} \binom{n}{k} \binom{n+k}{k-1}$$


and the **Narayana numbers**

$$\text{Nar}(n; k) = \frac{1}{n} \binom{n}{k} \binom{n}{k-1}.$$

The Classical Associahedron

- Example: Here are the f -vector and h -vector of $\text{Ass}(4)$.

				1			
			1		6		
		1		7		6	
	1		8		13		1
1		9		21		14	



The Rational Associahedron

Idea

Given $0 < x = a/(b - a)$ with $0 < a < b$ coprime, we will define a simplicial complex

$$\text{Ass}(x) = \text{Ass}(a, b)$$

whose maximal faces correspond to certain special dissections (“rational triangulations”) of a convex $(b + 1)$ -gon.

The Rational Associahedron

Idea


Given $0 < x = a/(b - a)$ with $0 < a < b$ coprime, we will define a simplicial complex

$$\text{Ass}(x) = \text{Ass}(a, b)$$

whose maximal faces correspond to certain special dissections (“rational triangulations”) of a convex $(b + 1)$ -gon.


To define a “rational triangulation” ...

- ▶ Start with a Dyck path. Here $(a, b) = (5, 8)$.




To define a “rational triangulation” ...

- ▶ Label the **columns** by $\{1, 2, \dots, b + 1\}$.




To define a “rational triangulation” ...

- ▶ Shoot **lasers** from the bottom left with **slope a/b** .




To define a “rational triangulation” ...

- ▶ Lift the lasers up.




To define a “rational triangulation” ...

- ▶ There you go!



To define a “rational triangulation” ...

- ▶ We have constructed $\text{Cat}(a, b)$ many “rational triangulations” of a convex $(b + 1)$ -gon, and **each of them has $a - 1$ chords**.



The Rational Associahedron

Definition

Given $0 < x = a/(b - a)$, let $\text{Ass}(x) = \text{Ass}(a, b)$ be the abstract simplicial complex whose maximal faces are the “rational triangulations”.

Geometric Realization

Note that $\text{Ass}(a, b)$ is a pure $(a - 1)$ -dimensional subcomplex of the $(b - 1)$ -dimensional polytope $\text{Ass}(b - 1)$.

The Rational Associahedron

Definition

Given $0 < x = a/(b - a)$, let $\text{Ass}(x) = \text{Ass}(a, b)$ be the abstract simplicial complex whose maximal faces are the “rational triangulations”.

Geometric Realization

Note that $\text{Ass}(a, b)$ is a pure $(a - 1)$ -dimensional subcomplex of the $(b - 1)$ -dimensional polytope $\text{Ass}(b - 1)$.

The Rational Associahedron

Theorems (with B. Rhoades and N. Williams)

- ▶ $\text{Ass}(n, n+1)$ is the **classical associahedron** $\text{Ass}(n)$.
- ▶ $\text{Ass}(n, (k-1)n+1)$ is the **generalized cluster complex** of Athanasiadis-Tzanaki and Fomin-Reading.
- ▶ $\text{Ass}(x)$ has $\text{Cat}(x)$ maximal faces and **Euler characteristic** $\text{Cat}'(x)$.
- ▶ $\text{Ass}(x)$ is **shellable** and hence homotopy equivalent to a wedge of $\text{Cat}'(x)$ many $(a-1)$ -dimensional spheres.
- ▶ $\text{Ass}(x)$ has **h -vector** $\text{Nar}(x; k) = \frac{1}{a} \binom{a}{k} \binom{b-1}{k-1}$.
- ▶ Hence its **f -vector** is given by the **rational Kirkman numbers**:

$$\text{Kirk}(x; k) := \frac{1}{a} \binom{a}{k} \binom{b+k-1}{k-1}.$$

The Rational Associahedron

Theorems (with B. Rhoades and N. Williams)

- ▶ $\text{Ass}(n, n + 1)$ is the **classical associahedron** $\text{Ass}(n)$.
- ▶ $\text{Ass}(n, (k - 1)n + 1)$ is the **generalized cluster complex** of Athanasiadis-Tzanaki and Fomin-Reading.
- ▶ $\text{Ass}(x)$ has $\text{Cat}(x)$ maximal faces and **Euler characteristic** $\text{Cat}'(x)$.
- ▶ $\text{Ass}(x)$ is **shellable** and hence homotopy equivalent to a wedge of $\text{Cat}'(x)$ many $(a - 1)$ -dimensional spheres.
- ▶ $\text{Ass}(x)$ has **h -vector** $\text{Nar}(x; k) = \frac{1}{a} \binom{a}{k} \binom{b-1}{k-1}$.
- ▶ Hence its **f -vector** is given by the **rational Kirkman numbers**:

$$\text{Kirk}(x; k) := \frac{1}{a} \binom{a}{k} \binom{b+k-1}{k-1}.$$

The Rational Associahedron

Theorems (with B. Rhoades and N. Williams)

- ▶ $\text{Ass}(n, n + 1)$ is the **classical associahedron** $\text{Ass}(n)$.
- ▶ $\text{Ass}(n, (k - 1)n + 1)$ is the **generalized cluster complex** of Athanasiadis-Tzanaki and Fomin-Reading.
- ▶ $\text{Ass}(x)$ has $\text{Cat}(x)$ maximal faces and **Euler characteristic** $\text{Cat}'(x)$.
- ▶ $\text{Ass}(x)$ is **shellable** and hence homotopy equivalent to a wedge of $\text{Cat}'(x)$ many $(a - 1)$ -dimensional spheres.
- ▶ $\text{Ass}(x)$ has **h -vector** $\text{Nar}(x; k) = \frac{1}{a} \binom{a}{k} \binom{b-1}{k-1}$.
- ▶ Hence its **f -vector** is given by the **rational Kirkman numbers**:

$$\text{Kirk}(x; k) := \frac{1}{a} \binom{a}{k} \binom{b+k-1}{k-1}.$$

The Rational Associahedron

Theorems (with B. Rhoades and N. Williams)

- ▶ $\text{Ass}(n, n + 1)$ is the **classical associahedron** $\text{Ass}(n)$.
- ▶ $\text{Ass}(n, (k - 1)n + 1)$ is the **generalized cluster complex** of Athanasiadis-Tzanaki and Fomin-Reading.
- ▶ $\text{Ass}(x)$ has $\text{Cat}(x)$ maximal faces and **Euler characteristic** $\text{Cat}'(x)$.
- ▶ $\text{Ass}(x)$ is **shellable** and hence homotopy equivalent to a wedge of $\text{Cat}'(x)$ many $(a - 1)$ -dimensional spheres.
- ▶ $\text{Ass}(x)$ has **h -vector** $\text{Nar}(x; k) = \frac{1}{a} \binom{a}{k} \binom{b-1}{k-1}$.
- ▶ Hence its **f -vector** is given by the **rational Kirkman numbers**:

$$\text{Kirk}(x; k) := \frac{1}{a} \binom{a}{k} \binom{b+k-1}{k-1}.$$

The Rational Associahedron

Theorems (with B. Rhoades and N. Williams)

- ▶ $\text{Ass}(n, n + 1)$ is the **classical associahedron** $\text{Ass}(n)$.
- ▶ $\text{Ass}(n, (k - 1)n + 1)$ is the **generalized cluster complex** of Athanasiadis-Tzanaki and Fomin-Reading.
- ▶ $\text{Ass}(x)$ has $\text{Cat}(x)$ maximal faces and **Euler characteristic** $\text{Cat}'(x)$.
- ▶ $\text{Ass}(x)$ is **shellable** and hence homotopy equivalent to a wedge of $\text{Cat}'(x)$ many $(a - 1)$ -dimensional spheres.
- ▶ $\text{Ass}(x)$ has **h -vector** $\text{Nar}(x; k) = \frac{1}{a} \binom{a}{k} \binom{b-1}{k-1}$.
- ▶ Hence its **f -vector** is given by the **rational Kirkman numbers**:

$$\text{Kirk}(x; k) := \frac{1}{a} \binom{a}{k} \binom{b+k-1}{k-1}.$$

The Rational Associahedron

Theorems (with B. Rhoades and N. Williams)

- ▶ $\text{Ass}(n, n + 1)$ is the **classical associahedron** $\text{Ass}(n)$.
- ▶ $\text{Ass}(n, (k - 1)n + 1)$ is the **generalized cluster complex** of Athanasiadis-Tzanaki and Fomin-Reading.
- ▶ $\text{Ass}(x)$ has $\text{Cat}(x)$ maximal faces and **Euler characteristic** $\text{Cat}'(x)$.
- ▶ $\text{Ass}(x)$ is **shellable** and hence homotopy equivalent to a wedge of $\text{Cat}'(x)$ many $(a - 1)$ -dimensional spheres.
- ▶ $\text{Ass}(x)$ has **h -vector** $\text{Nar}(x; k) = \frac{1}{a} \binom{a}{k} \binom{b-1}{k-1}$.
- ▶ Hence its **f -vector** is given by the **rational Kirkman numbers**:

$$\text{Kirk}(x; k) := \frac{1}{a} \binom{a}{k} \binom{b+k-1}{k-1}.$$

The Rational Associahedron

Theorems (with B. Rhoades and N. Williams)

- ▶ $\text{Ass}(n, n + 1)$ is the **classical associahedron** $\text{Ass}(n)$.
- ▶ $\text{Ass}(n, (k - 1)n + 1)$ is the **generalized cluster complex** of Athanasiadis-Tzanaki and Fomin-Reading.
- ▶ $\text{Ass}(x)$ has $\text{Cat}(x)$ maximal faces and **Euler characteristic** $\text{Cat}'(x)$.
- ▶ $\text{Ass}(x)$ is **shellable** and hence homotopy equivalent to a wedge of $\text{Cat}'(x)$ many $(a - 1)$ -dimensional spheres.
- ▶ $\text{Ass}(x)$ has **h -vector** $\text{Nar}(x; k) = \frac{1}{a} \binom{a}{k} \binom{b-1}{k-1}$.
- ▶ Hence its **f -vector** is given by the **rational Kirkman numbers**:

$$\text{Kirk}(x; k) := \frac{1}{a} \binom{a}{k} \binom{b+k-1}{k-1}.$$

The Rational Associahedron

Observation

Given $0 < x = a/(b - a)$ with $0 < a < b$ coprime, note that $\text{Ass}(x) = \text{Ass}(a, b)$ and $\text{Ass}(1/x) = \text{Ass}(b - a, b)$ are both subcomplexes of the polytope $\text{Ass}(b - 1) = \text{Ass}(b - 1, b)$.

Question

How do $\text{Ass}(x)$ and $\text{Ass}(1/x)$ fit together?

The Rational Associahedron

Observation


Given $0 < x = a/(b - a)$ with $0 < a < b$ coprime, note that $\text{Ass}(x) = \text{Ass}(a, b)$ and $\text{Ass}(1/x) = \text{Ass}(b - a, b)$ are both subcomplexes of the polytope $\text{Ass}(b - 1) = \text{Ass}(b - 1, b)$.

Question

How do $\text{Ass}(x)$ and $\text{Ass}(1/x)$ fit together?

The Rational Associahedron

- ▶ Example: Here are subcomplexes $\text{Ass}(2, 5)$ and $\text{Ass}(3, 5)$ in $\text{Ass}(4)$.



Rational Duality?

Observation

Note that $\text{Ass}(b-1)$ has this many vertices:

$$\binom{b+1}{2} - (b+1) = \frac{(b+1)b}{2} - \frac{2(b+1)}{2} = \frac{(b-2)(b+1)}{2}.$$

The subcomplexes $\text{Ass}(a, b)$ and $\text{Ass}(b-a, b)$ **bipartition** the vertices:

$$\frac{(a-1)(b+1)}{2} + \frac{(b-a-1)(b+1)}{2} = \frac{(b-2)(b+1)}{2}.$$

Rational Duality = Alexander Duality

Conjecture (with B. Rhoades and N. Williams)

We know that $\text{Ass}(a, b)$ and $\text{Ass}(b - a, b)$ have the same number of homotopy spheres (of complementary dimensions) because

$$\text{Cat}'(a, b) = \text{Cat}'(b - a, b).$$

*We conjecture that the homotopy spheres are “intertwined” in a nice way. Formally, we conjecture that $\text{Ass}(a, b)$ and $\text{Ass}(b - a, b)$ are **Alexander dual** as subcomplexes of $\text{Ass}(b - 1)$.*

Theorem (B. Rhoades)

The conjecture is true.

Rational Duality = Alexander Duality

Conjecture (with B. Rhoades and N. Williams)

We know that $\text{Ass}(a, b)$ and $\text{Ass}(b - a, b)$ have the same number of homotopy spheres (of complementary dimensions) because

$$\text{Cat}'(a, b) = \text{Cat}'(b - a, b).$$

*We conjecture that the homotopy spheres are “intertwined” in a nice way. Formally, we conjecture that $\text{Ass}(a, b)$ and $\text{Ass}(b - a, b)$ are **Alexander dual** as subcomplexes of $\text{Ass}(b - 1)$.*

Theorem (B. Rhoades)

The conjecture is true.

Rational Duality = Alexander Duality

Conjecture (with B. Rhoades and N. Williams)

We know that $\text{Ass}(a, b)$ and $\text{Ass}(b - a, b)$ have the same number of homotopy spheres (of complementary dimensions) because

$$\text{Cat}'(a, b) = \text{Cat}'(b - a, b).$$

*We conjecture that the homotopy spheres are “intertwined” in a nice way. Formally, we conjecture that $\text{Ass}(a, b)$ and $\text{Ass}(b - a, b)$ are **Alexander dual** as subcomplexes of $\text{Ass}(b - 1)$.*

Theorem (B. Rhoades)

The conjecture is true.

Euclidean Algorithm = ?

Definition

Given $0 < a < b$ coprime, if we define

$$\text{Ass}'(a, b) := \begin{cases} \text{Ass}(a, b - a) & \text{for } a < (b - a) \\ \text{Ass}(b - a, a) & \text{for } (b - a) < a \end{cases}$$

then

homotopy spheres $\text{Ass}(a, b) = \#$ **maximal faces** $\text{Ass}'(a, b)$.

Question

What does the following mean?

$$\text{Ass}(a, b) \mapsto \text{Ass}'(a, b) \mapsto \text{Ass}''(a, b) \mapsto \cdots \mapsto \mathbf{a \ point}$$

Euclidean Algorithm = ?

Definition

Given $0 < a < b$ coprime, if we define

$$\text{Ass}'(a, b) := \begin{cases} \text{Ass}(a, b - a) & \text{for } a < (b - a) \\ \text{Ass}(b - a, a) & \text{for } (b - a) < a \end{cases}$$

then

homotopy spheres $\text{Ass}(a, b) = \text{\# maximal faces}$ $\text{Ass}'(a, b)$.

Question

What does the following mean?

$$\text{Ass}(a, b) \mapsto \text{Ass}'(a, b) \mapsto \text{Ass}''(a, b) \mapsto \cdots \mapsto \text{a point}$$

Euclidean Algorithm = ?

Definition

Given $0 < a < b$ coprime, if we define

$$\text{Ass}'(a, b) := \begin{cases} \text{Ass}(a, b - a) & \text{for } a < (b - a) \\ \text{Ass}(b - a, a) & \text{for } (b - a) < a \end{cases}$$

then

homotopy spheres $\text{Ass}(a, b) = \text{\# maximal faces}$ $\text{Ass}'(a, b)$.

Question

What does the following mean?


$$\text{Ass}(a, b) \mapsto \text{Ass}'(a, b) \mapsto \text{Ass}''(a, b) \mapsto \cdots \mapsto \text{a point}$$

What is a Parking Function?

The Rational Parking Space

Definition

- ▶ Label the up-steps by $\{1, 2, \dots, a\}$, increasing up columns.




- ▶ Call this a **parking function**.
- ▶ Let $\text{PF}(x) = \text{PF}(a, b)$ denote the set of parking functions.
- ▶ Classical form (z_1, z_2, \dots, z_a) has label z_i in column i .
- ▶ Example: $(3, 1, 4, 4, 1)$

The Rational Parking Space

Definition

- ▶ Label the up-steps by $\{1, 2, \dots, a\}$, increasing up columns.




- ▶ Call this a **parking function**.
- ▶ Let $\text{PF}(x) = \text{PF}(a, b)$ denote the set of parking functions.
- ▶ **Classical form** (z_1, z_2, \dots, z_a) has label z_i in column i .
- ▶ Example: $(3, 1, 4, 4, 1)$

The Rational Parking Space

Definition

- ▶ Label the up-steps by $\{1, 2, \dots, a\}$, increasing up columns.




- ▶ Call this a **parking function**.
- ▶ Let $PF(x) = PF(a, b)$ denote the set of parking functions.
- ▶ Classical form (z_1, z_2, \dots, z_a) has label z_i in column i .
- ▶ Example: $(3, 1, 4, 4, 1)$

The Rational Parking Space

Definition

- ▶ Label the up-steps by $\{1, 2, \dots, a\}$, increasing up columns.




- ▶ Call this a **parking function**.
- ▶ Let $\text{PF}(x) = \text{PF}(a, b)$ denote the set of parking functions.
- ▶ **Classical form** (z_1, z_2, \dots, z_a) has label z_i in column i .
- ▶ Example: $(3, 1, 4, 4, 1)$

The Rational Parking Space

Definition

- ▶ The symmetric group \mathfrak{S}_a acts on classical forms.




- ▶ Example: $(3, 1, 4, 4, 1)$ versus $(3, 1, 1, 4, 4)$
- ▶ By abuse, let $\text{PF}(x) = \text{PF}(a, b)$ denote this representation of \mathfrak{S}_a .
- ▶ Call it the **rational parking space**.

The Rational Parking Space

Definition

- ▶ The symmetric group \mathfrak{S}_a acts on classical forms.




- ▶ Example: $(3, 1, 4, 4, 1)$ versus $(3, 1, 1, 4, 4)$
 - ▶ By abuse, let $\text{PF}(x) = \text{PF}(a, b)$ denote this representation of \mathfrak{S}_a .
 - ▶ Call it the **rational parking space**.

The Rational Parking Space

Definition

- ▶ The symmetric group \mathfrak{S}_a acts on classical forms.




- ▶ Example: $(3, 1, 4, 4, 1)$ versus $(3, 1, 1, 4, 4)$
- ▶ By abuse, let $\text{PF}(x) = \text{PF}(a, b)$ denote this representation of \mathfrak{S}_a .
- ▶ Call it the **rational parking space**.

The Rational Parking Space

Definition

- ▶ The symmetric group \mathfrak{S}_a acts on classical forms.




- ▶ Example: $(3, 1, 4, 4, 1)$ versus $(3, 1, 1, 4, 4)$
- ▶ By abuse, let $\text{PF}(x) = \text{PF}(a, b)$ denote this representation of \mathfrak{S}_a .
- ▶ Call it the **rational parking space**.

The Rational Parking Space

Definition

- ▶ The symmetric group \mathfrak{S}_a acts on classical forms.



- ▶ Example: $(3, 1, 4, 4, 1)$ versus $(3, 1, 1, 4, 4)$
- ▶ By abuse, let $\text{PF}(x) = \text{PF}(a, b)$ denote this representation of \mathfrak{S}_a .
- ▶ Call it the **rational parking space**.

The Rational Parking Space

Theorems (with N. Loehr and G. Warrington)

- ▶ The dimension of $\text{PF}(a, b)$ is b^{a-1} .
- ▶ The **complete homogeneous expansion** is

$$\text{PF}(a, b) = \sum_{r \vdash a} \frac{1}{b} \binom{b}{r_0, r_1, \dots, r_a} h_r,$$

where the sum is over $r = 0^{r_0} 1^{r_1} \dots a^{r_a} \vdash a$ with $\sum_i r_i = b$.

- ▶ Note that this is the same as

$$\text{PF}(a, b) = \sum_{r \vdash a} \frac{1}{b} m_r(1^b) h_r.$$

The Rational Parking Space

Theorems (with N. Loehr and G. Warrington)

► The dimension of $\text{PF}(a, b)$ is b^{a-1} .

► The complete homogeneous expansion is

$$\text{PF}(a, b) = \sum_{\mathbf{r} \vdash a} \frac{1}{b} \binom{b}{r_0, r_1, \dots, r_a} h_{\mathbf{r}},$$

where the sum is over $\mathbf{r} = 0^{r_0} 1^{r_1} \dots a^{r_a} \vdash a$ with $\sum_i r_i = b$.

► Note that this is the same as

$$\text{PF}(a, b) = \sum_{\mathbf{r} \vdash a} \frac{1}{b} m_{\mathbf{r}}(1^b) h_{\mathbf{r}}.$$

The Rational Parking Space

Theorems (with N. Loehr and G. Warrington)

- ▶ The dimension of $\text{PF}(a, b)$ is b^{a-1} .
- ▶ The **complete homogeneous expansion** is

$$\text{PF}(a, b) = \sum_{\mathbf{r} \vdash a} \frac{1}{b} \binom{b}{r_0, r_1, \dots, r_a} h_{\mathbf{r}},$$

where the sum is over $\mathbf{r} = 0^{r_0} 1^{r_1} \dots a^{r_a} \vdash a$ with $\sum_i r_i = b$.

- ▶ Note that this is the same as

$$\text{PF}(a, b) = \sum_{\mathbf{r} \vdash a} \frac{1}{b} m_{\mathbf{r}}(1^b) h_{\mathbf{r}}.$$

The Rational Parking Space

Theorems (with N. Loehr and G. Warrington)

- ▶ The dimension of $\text{PF}(a, b)$ is b^{a-1} .
- ▶ The **complete homogeneous expansion** is

$$\text{PF}(a, b) = \sum_{\mathbf{r} \vdash a} \frac{1}{b} \binom{b}{r_0, r_1, \dots, r_a} h_{\mathbf{r}},$$

where the sum is over $\mathbf{r} = 0^{r_0} 1^{r_1} \dots a^{r_a} \vdash a$ with $\sum_i r_i = b$.

- ▶ Note that this is the same as

$$\text{PF}(a, b) = \sum_{\mathbf{r} \vdash a} \frac{1}{b} m_{\mathbf{r}}(1^b) h_{\mathbf{r}}.$$

A Few Facts

Theorems (with N. Loehr and G. Warrington)

Then using the Cauchy product identity we get...

- ▶ The **power sum expansion** is

$$\text{PF}(a, b) = \sum_{r \vdash a} b^{\ell(r)-1} \frac{p_r}{z_r}$$

i.e. the # of parking functions fixed by $\sigma \in \mathfrak{S}_a$ is $b^{\#\text{cycles}(\sigma)-1}$.

- ▶ The **Schur expansion** is

$$\text{PF}(a, b) = \sum_{r \vdash a} \frac{1}{b} s_r(1^b) s_r.$$

A Few Facts

Theorems (with N. Loehr and G. Warrington)

Then using the Cauchy product identity we get...

- ▶ The **power sum expansion** is

$$\text{PF}(a, b) = \sum_{r \vdash a} b^{\ell(r)-1} \frac{p_r}{z_r}$$

i.e. the # of parking functions fixed by $\sigma \in \mathfrak{S}_a$ is $b^{\#\text{cycles}(\sigma)-1}$.

- ▶ The **Schur expansion** is

$$\text{PF}(a, b) = \sum_{r \vdash a} \frac{1}{b} s_r(1^b) s_r.$$

A Few Facts

Theorems (with N. Loehr and G. Warrington)

Then using the Cauchy product identity we get...

- ▶ The **power sum expansion** is

$$\text{PF}(a, b) = \sum_{r \vdash a} b^{\ell(r)-1} \frac{p_r}{z_r}$$

i.e. the # of parking functions fixed by $\sigma \in \mathfrak{S}_a$ is $b^{\#\text{cycles}(\sigma)-1}$.

- ▶ The **Schur expansion** is

$$\text{PF}(a, b) = \sum_{r \vdash a} \frac{1}{b} s_r(1^b) s_r.$$

A Few Facts

Observation/Definition

The multiplicities of the **hook Schur functions** $s[k+1, 1^{a-k-1}]$ in $\text{PF}(a, b)$ are given by the **rational Schröder numbers**:

$$\text{Schrö}(a, b; k) := \frac{1}{b} s_{[k+1, 1^{a-k-1}]}(1^b) = \frac{1}{b} \binom{a-1}{k} \binom{b+k}{a}.$$

Special Cases:

- ▶ Trivial character: $\text{Schrö}(a, b; a-1) = \text{Cat}(a, b)$.
- ▶ Smallest k that occurs is $k = \max\{0, a-b\}$, in which case

$$\text{Schrö}(a, b; k) = \text{Cat}'(a, b).$$

- ▶ Hence $\text{Schrö}(x; k)$ interpolates between $\text{Cat}(x)$ and $\text{Cat}'(x)$.

A Few Facts

Observation/Definition

The multiplicities of the **hook Schur functions** $s[k+1, 1^{a-k-1}]$ in $\text{PF}(a, b)$ are given by the **rational Schröder numbers**:

$$\text{Schrö}(a, b; k) := \frac{1}{b} s_{[k+1, 1^{a-k-1}]}(1^b) = \frac{1}{b} \binom{a-1}{k} \binom{b+k}{a}.$$

Special Cases:

- ▶ Trivial character: $\text{Schrö}(a, b; a-1) = \text{Cat}(a, b)$.
- ▶ Smallest k that occurs is $k = \max\{0, a-b\}$, in which case

$$\text{Schrö}(a, b; k) = \text{Cat}'(a, b).$$

- ▶ Hence $\text{Schrö}(x; k)$ interpolates between $\text{Cat}(x)$ and $\text{Cat}'(x)$.

A Few Facts

Observation/Definition

The multiplicities of the **hook Schur functions** $s[k+1, 1^{a-k-1}]$ in $\text{PF}(a, b)$ are given by the **rational Schröder numbers**:

$$\text{Schrö}(a, b; k) := \frac{1}{b} s_{[k+1, 1^{a-k-1}]}(1^b) = \frac{1}{b} \binom{a-1}{k} \binom{b+k}{a}.$$

Special Cases:

- ▶ Trivial character: $\text{Schrö}(a, b; a-1) = \text{Cat}(a, b)$.
- ▶ Smallest k that occurs is $k = \max\{0, a-b\}$, in which case

$$\text{Schrö}(a, b; k) = \text{Cat}'(a, b).$$

- ▶ Hence $\text{Schrö}(x; k)$ interpolates between $\text{Cat}(x)$ and $\text{Cat}'(x)$.

A Few Facts

Observation/Definition

The multiplicities of the **hook Schur functions** $s[k+1, 1^{a-k-1}]$ in $\text{PF}(a, b)$ are given by the **rational Schröder numbers**:

$$\text{Schrö}(a, b; k) := \frac{1}{b} s_{[k+1, 1^{a-k-1}]}(1^b) = \frac{1}{b} \binom{a-1}{k} \binom{b+k}{a}.$$

Special Cases:

- ▶ Trivial character: $\text{Schrö}(a, b; a-1) = \text{Cat}(a, b)$.
- ▶ Smallest k that occurs is $k = \max\{0, a-b\}$, in which case

$$\text{Schrö}(a, b; k) = \text{Cat}'(a, b).$$

- ▶ Hence $\text{Schrö}(x; k)$ interpolates between $\text{Cat}(x)$ and $\text{Cat}'(x)$.

A Few Facts

Observation/Definition

The multiplicities of the **hook Schur functions** $s[k+1, 1^{a-k-1}]$ in $\text{PF}(a, b)$ are given by the **rational Schröder numbers**:

$$\text{Schrö}(a, b; k) := \frac{1}{b} s_{[k+1, 1^{a-k-1}]}(1^b) = \frac{1}{b} \binom{a-1}{k} \binom{b+k}{a}.$$

Special Cases:

- ▶ Trivial character: $\text{Schrö}(a, b; a-1) = \text{Cat}(a, b)$.
- ▶ Smallest k that occurs is $k = \max\{0, a-b\}$, in which case

$$\text{Schrö}(a, b; k) = \text{Cat}'(a, b).$$

- ▶ Hence $\text{Schrö}(x; k)$ interpolates between $\text{Cat}(x)$ and $\text{Cat}'(x)$.

What does switching $a \leftrightarrow b$ mean?

Problem

Given a, b coprime we have an \mathfrak{S}_a -module $\text{PF}(a, b)$ of dimension b^{a-1} and an \mathfrak{S}_b -module $\text{PF}(b, a)$ of dimension a^{b-1} .

- ▶ What is the relationship between $\text{PF}(a, b)$ and $\text{PF}(b, a)$?
- ▶ Note that hook multiplicities are the same:

$$\text{Schrö}(a, b; k) = \text{Schrö}(b, a; k + b - a).$$

- ▶ See: *E. Gorsky, "Arc spaces and DAHA representations", (2011)*

What does switching $a \leftrightarrow b$ mean?

Problem

Given a, b coprime we have an \mathfrak{S}_a -module $\text{PF}(a, b)$ of dimension b^{a-1} and an \mathfrak{S}_b -module $\text{PF}(b, a)$ of dimension a^{b-1} .

- ▶ What is the relationship between $\text{PF}(a, b)$ and $\text{PF}(b, a)$?
- ▶ Note that hook multiplicities are the same:

$$\text{Schrö}(a, b; k) = \text{Schrö}(b, a; k + b - a).$$

- ▶ See: *E. Gorsky, "Arc spaces and DAHA representations", (2011)*

What does switching $a \leftrightarrow b$ mean?

Problem

Given a, b coprime we have an \mathfrak{S}_a -module $\text{PF}(a, b)$ of dimension b^{a-1} and an \mathfrak{S}_b -module $\text{PF}(b, a)$ of dimension a^{b-1} .

- ▶ What is the relationship between $\text{PF}(a, b)$ and $\text{PF}(b, a)$?
- ▶ Note that hook multiplicities are the same:

$$\text{Schrö}(a, b; k) = \text{Schrö}(b, a; k + b - a).$$

- ▶ See: *E. Gorsky, "Arc spaces and DAHA representations", (2011)*

What does switching $a \leftrightarrow b$ mean?

Problem

Given a, b coprime we have an \mathfrak{S}_a -module $\text{PF}(a, b)$ of dimension b^{a-1} and an \mathfrak{S}_b -module $\text{PF}(b, a)$ of dimension a^{b-1} .

- ▶ What is the relationship between $\text{PF}(a, b)$ and $\text{PF}(b, a)$?
- ▶ Note that hook multiplicities are the same:

$$\text{Schrö}(a, b; k) = \text{Schrö}(b, a; k + b - a).$$

- ▶ See: *E. Gorsky, "Arc spaces and DAHA representations", (2011)*

Summary of Catalan Numerology

- ▶ The Kirkman/Narayana/Schröder numbers are equivalent. They contain information about rank. ($1 < k < a - 1$)

$$\left. \begin{aligned} \text{Kirk}(x; k) &= \frac{1}{a} \binom{a}{k} \binom{b+k-1}{k-1} \\ \text{Nar}(x; k) &= \frac{1}{a} \binom{a}{k} \binom{b-1}{k-1} \\ \text{Schrö}(x; k) &= \frac{1}{b} \binom{a-1}{k} \binom{b+k}{a} \end{aligned} \right\} \begin{array}{l} f\text{-vector} \\ h\text{-vector} \\ \text{"dual" } f\text{-vector} \end{array}$$

- ▶ The Kreweras numbers are more refined. They contain parabolic information. ($\mathbf{r} \vdash a$)

$$\text{Krew}(x; \mathbf{r}) = \frac{1}{b} \binom{b}{r_0, r_1, \dots, r_a}$$

Summary of Catalan Numerology

- ▶ The Kirkman/Narayana/Schröder numbers are equivalent. They contain information about rank. ($1 < k < a - 1$)

$$\left. \begin{aligned} \text{Kirk}(x; k) &= \frac{1}{a} \binom{a}{k} \binom{b+k-1}{k-1} \\ \text{Nar}(x; k) &= \frac{1}{a} \binom{a}{k} \binom{b-1}{k-1} \\ \text{Schrö}(x; k) &= \frac{1}{b} \binom{a-1}{k} \binom{b+k}{a} \end{aligned} \right\} \begin{array}{l} f\text{-vector} \\ h\text{-vector} \\ \text{"dual" } f\text{-vector} \end{array}$$

- ▶ The Kreweras numbers are more refined. They contain parabolic information. ($\mathbf{r} \vdash a$)

$$\text{Krew}(x; \mathbf{r}) = \frac{1}{b} \binom{b}{r_0, r_1, \dots, r_a}$$

But what about q and t ?

Tease

There **exists** a bigraded version $\text{PF}_{q,t}(a, b)$. Here is the coefficient of the (non-hook) Schur function $s[2, 2, 1]$ in $\text{PF}_{q,t}(5, 8)$:

$$\begin{pmatrix} & & & & & & 1 & 1 & 1 & 1 & 1 \\ & & & & & 1 & 3 & 4 & 3 & 2 & 1 \\ & & & 2 & 6 & 6 & 4 & 2 & 1 & & \\ & & 2 & 7 & 7 & 4 & 2 & 1 & & & \\ & 1 & 6 & 7 & 4 & 2 & 1 & & & & \\ & 3 & 6 & 4 & 2 & 1 & & & & & \\ 1 & 4 & 4 & 2 & 1 & & & & & & \\ 1 & 3 & 2 & 1 & & & & & & & \\ 1 & 2 & 1 & & & & & & & & \\ 1 & 1 & & & & & & & & & \\ 1 & & & & & & & & & & \end{pmatrix}$$

Vielen Dank!

