

STABLE PERIODIC OSCILLATIONS IN A TWO-STAGE CANCER MODEL OF TUMOR AND IMMUNE SYSTEM INTERACTIONS

DAN LIU

Department of Mathematics, Xidian University
Xi'an, Shanxi 710071, China

SHIGUI RUAN

Department of Mathematics, University of Miami,
Coral Gables, FL 33124, USA

DEMING ZHU

Department of Mathematics, East China Normal University,
Shanghai 200062, P. R. China

(Communicated by Stephen Gourley)

ABSTRACT. This paper presents qualitative and bifurcation analysis near the degenerate equilibrium in a two-stage cancer model of interactions between lymphocyte cells and solid tumor and contributes to a better understanding of the dynamics of tumor and immune system interactions. We first establish the existence of Hopf bifurcation in the 3-dimensional cancer model and rule out the occurrence of the degenerate Hopf bifurcation. Then a general Hopf bifurcation formula is applied to determine the stability of the limit cycle bifurcated from the interior equilibrium. Sufficient conditions on the existence of stable periodic oscillations of tumor levels are obtained for the two-stage cancer model. Numerical simulations are presented to illustrate the existence of stable periodic oscillations with reasonable parameters and demonstrate the phenomenon of long-term tumor relapse in the model.

1. Introduction. According to the study in Boyle et al. [3], every year millions of people suffer with cancer and die from this disease throughout the world. In the new century cancer still remains one of the most dangerous killers of humankind. In order to study the progress of cancer and seek effective treatment strategies, researchers have proposed comprehensive approaches, including biological, computational and mathematical methods, to study the disease. An important aspect in studying cancer is to understand the interactions between the immune system and tumor cells. In order to simulate the host's own immune response to destroy and eliminate tumor cells, various types of mathematical models have been proposed, see Albert et al. [1], de Pillis et al. [6], d'Onofrio [7], Kirschner and Panetta [15], Kirschner and Tsyvintsev [16], Kuznetsov et al. [17], Lejeune et al. [19], Liu et al. [22], and Swan [30]. We refer to a recent survey by Eftimie et al. [9] and the references cited

2000 *Mathematics Subject Classification.* Primary: 34C23, 34C60; Secondary: 37G10.

Key words and phrases. Hopf bifurcation, equilibrium, oscillation, tumor, lymphocyte.

The first author was supported by NSFC grant 10801051. The second author was partially supported by NSF grant DMS-1022728.

therein for more references on modeling tumor and immune system interactions, in particular using ordinary differential equations.

In this work we are interested in studying mathematical models of carcinogenesis in which the tumor cells elicit an immune response. Based on some reasonable hypotheses, the following model

$$\begin{aligned}\frac{dL}{dt} &= -\lambda_1 L + \alpha'_1 \bar{C} L \left(1 - \frac{L}{L_c}\right), \\ \frac{dC}{dt} &= \lambda_2 C_f - \alpha'_2 \bar{C} L\end{aligned}\tag{1}$$

was proposed by DeLisi and Rescigno [5] to qualitatively estimate the function of the immune surveillance (Lefever and Garay [18]), where $L(t)$ and $C(t)$ denote respectively the number of free lymphocytes and the total number of tumor cells at time t , \bar{C} and C_f are the total number of free tumor cells and the number of free cells on a tumor surface (i.e., not bounded by lymphocytes), respectively. The values of λ_1 and α'_1 may be dependent on environmental stimuli and the type of tissues where the tumor is growing. The last factor in the first equation $1 - \frac{L}{L_c}$ is a saturation term which restricts the maximum number L_c of lymphocytes in the system. If the relationship between free and bounded lymphocytes is assumed to be equilibrium controlled, with K the equilibrium constant for lymphocyte and tumor cell interaction, and the tumor is assumed to be spherical, then DeLisi and Rescigno obtained that $C_f = C - gKLC^{2/3}/(1+KL)$, $\bar{C} = gC^{2/3}/(1+KL)$, where $g > 0$ is a constant. DeLisi and Rescigno [5] analyzed the qualitative behavior of the solutions of model (1) except near the degenerate equilibrium and compared briefly with the results of the transplantation experiments. Recently, Lin [20] studied the existence of the solutions and the stability of the steady states of model (1). The dynamical behavior and bifurcations near the degenerate equilibrium in model (1) was studied by Liu et al. [21].

System (1) describes a mathematical model of the interactions between the immune system and solid tumor. This model is consistent with the experimental observations in many conclusions, however, the development of de novo tumor is unable to be predicted. To avoid this difficulty, Rescigno and DeLisi [26] made slight but more realistic changes on the model by requiring that lymphocytes go through two stages of development, namely immature and mature. They claimed that only lymphocytes in the second stage are effective in killing tumor cells, which were referred to as activated lymphocytes. We would like to mention that other researchers have also considered different stages of immune cells or different immune cells in modeling tumor and immune system interactions. For example, in modeling cytotoxic reactions mediated by the response of immune cells to solid tumors, Lejeune et al. [19] argued that lysis and binding are independent processes and modeled free and bound immune cells separately. De Pillis et al. [6] proposed a model of cell-mediated immune response to tumor growth and considered the interaction of the natural killer cells and the CD8⁺ T cells with tumor cells separately.

Following Rescigno and DeLisi [26], we denote

$L_i(t)$ – the number of lymphocytes in stage i ($i = 1, 2$),

$C(t)$ – the total number of cells in the tumor,

$C_f(t)$ – the total number of free (i.e., unbounded) cancer cells,

and make the following assumptions:

- (a) The tumor is spherical at all times and its geometry is protective so that only cells on the surface of the tumor are vulnerable to attack.

- (b) The death rate of the stage-2 lymphocytes is a first-order process with constant α_3 .
- (c) Stage-1 lymphocytes are transformed into stage 2 at a constant rate λ_1 .
- (d) Stage-1 lymphocytes are produced at a fixed rate $\lambda_1 L_0$ (in the absence of tumor cells) plus a rate proportional to the product of the number of free tumor cells and stage-2 lymphocytes.
- (e) In the absence of lymphocytes, the reproduction rate of the tumor cells is at a constant rate λ_2 .
- (f) The killing rate of the tumor cells by lymphocytes is proportional to the frequency of interaction with stage-2 lymphocytes with constant α'_2 .
- (g) There is an equilibrium relation between free tumor cells and stage-2 lymphocytes.
- (h) The progeny of L_2 must first pass through stage one before becoming mature lymphocytes to approximate some biological delay in the formation of L_2 by tumor cell stimulation.

Now the interactions between the tumor cells and lymphocytes are described a system of three ordinary differential equations (see [26]):

$$\begin{aligned} \frac{dL_1}{dt} &= -\lambda_1(L_1 - L_0) + \alpha'_1 C_f L_2 \exp(-L_2/L_c), \\ \frac{dL_2}{dt} &= \lambda_1 L_1 - \alpha_3 L_2, \\ \frac{dC}{dt} &= \lambda_2 C_f - \alpha'_2 C_f L_2, \end{aligned} \tag{2}$$

where $\alpha'_1 \exp(-L_2/L_c)$ represents the saturation of the system which restricts the maximum number L_c of lymphocytes, $(C - C_f)/C_f L_2$ is a constant K under the assumption that the relation between stage-2 lymphocytes and free tumor cells is equilibrium controlled. Therefore, $C_f = C/(1 + KL_2)$ and (2) is written as

$$\begin{aligned} \frac{dL_1}{dt} &= -\lambda_1(L_1 - L_0) + \alpha'_1 \frac{L_2 C}{1 + KL_2} \exp(-\frac{L_2}{L_c}), \\ \frac{dL_2}{dt} &= \lambda_1 L_1 - \alpha_3 L_2, \\ \frac{dC}{dt} &= \frac{\lambda_2 - \alpha'_2 L_2}{1 + KL_2} C, \end{aligned} \tag{3}$$

Some potential oscillation phenomena were observed by Rescigno and DeLisi [26] in the development of cancer and lymphocyte cells. In this paper we study the nonlinear dynamics of system (3) and prove that there is a stable periodic solution bifurcated from the interior degenerate equilibrium under certain assumptions of the parameters, which means that all the trajectories in the neighborhood of this equilibrium spiral towards the bifurcated limit cycle as time increases. More precisely, by applying the Hopf bifurcation theorem (Zhang [32]) for 3-dimensional systems, we first discuss the Hopf bifurcation in model (3) and obtain the nonexistence of degenerate Hopf bifurcation. The noticeable point lies in the determination of conditions on the stability of the bifurcated periodic orbit so that this cancer model exhibits stable periodic oscillations between the solid tumor cells and lymphocyte cells. Therefore, the tumor levels will oscillate inescapably once the initial values of tumor cells and lymphocyte cells are close enough to the degenerate equilibrium. This oscillatory phenomenon of tumor levels has been observed clinically and is known as Jeff's Phenomenon (Thomlinson [31]). The theoretical results in this work will be helpful for the further study of the dynamical behaviors

in the mathematical models of carcinogenesis and for the better understanding of the development of cancer when the tumor cells elicit an immune response.

The paper is organized as follows. Section 2 provides the qualitative analysis, the existence of Hopf bifurcation and the nonexistence of degenerate Hopf bifurcation. In section 3, an applicable transformation of a 3×3 Jacobian matrix is given. Section 4 discusses the stability of the bifurcated periodic orbit and presents the sufficient conditions on the stable case. Numerical simulations supporting the theoretical results of section 4 are presented in section 5. Section 6 provides a brief discussion on this work.

2. Hopf bifurcation.

For system (3), we make the following transformations: $x = K(L_1 - L_0)$, $y = KL_2$, $z = KC$, $\alpha_1 = \alpha'_1/K$, $\alpha_2 = \alpha'_2/K$, $x_0 = KL_0$, $y_c = KL_c$, then it is changed into

$$\begin{aligned} \frac{dx}{dt} &= -\lambda_1 x + \alpha_1 \frac{yz}{1+y} \exp(-\frac{y}{y_c}) \triangleq f(x, y, z), \\ \frac{dy}{dt} &= \lambda_1(x + x_0) - \alpha_3 y \triangleq g(x, y, z), \\ \frac{dz}{dt} &= \frac{\lambda_2 - \alpha_2 y}{1+y} z \triangleq h(x, y, z). \end{aligned} \tag{4}$$

The above system has two possible fixed points: $A(x_A, y_A, z_A) = (0, \frac{\lambda_1 x_0}{\alpha_3}, 0)$ and $B(x_B, y_B, z_B) = (\frac{\alpha_3 \lambda_2}{\alpha_2 \lambda_1} - x_0, \frac{\lambda_2}{\alpha_2}, \frac{\lambda_1 x_B (1+y_B)}{\alpha_1 y_B} \exp(\frac{y_B}{y_c}))$. The Jacobian matrix of the linearized system of (4) is

$$J(x, y, z) = \begin{pmatrix} -\lambda_1 & \alpha_1 z \frac{1-y/y_c - y^2/y_c}{(1+y)^2} \exp(-\frac{y}{y_c}) & \frac{\alpha_1 y}{1+y} \exp(-\frac{y}{y_c}) \\ \lambda_1 & -\alpha_3 & 0 \\ 0 & -\frac{(\alpha_2 + \lambda_2)z}{(1+y)^2} & \frac{\lambda_2 - \alpha_2 y}{1+y} \end{pmatrix}. \tag{5}$$

From the biological point of view, the state space of system (4) is the non-negative octant

$$\mathbb{R}_+^3 = \{X = (x, y, z)^T \in \mathbb{R}^3 : x \geq 0, y \geq 0, z \geq 0\},$$

where T means the transpose of a matrix or a vector throughout this paper. Denote the positive open subset of \mathbb{R}_+^3 as

$$\mathbb{R}_p^3 = \{X = (x, y, z)^T \in \mathbb{R}^3 : x > 0, y > 0, z > 0\}.$$

If we set $R_0 = \frac{\alpha_3 \lambda_2}{\alpha_2 \lambda_1 x_0}$, then the following conclusion can be obtained directly from [26].

Lemma 2.1. *If $R_0 < 1$, then for any solution $(x(t), y(t), z(t))$ of (4), we have that*

$$\liminf_{t \rightarrow +\infty} x(t) \geq 0, \liminf_{t \rightarrow +\infty} y(t) \geq y_A, \liminf_{t \rightarrow +\infty} z'(t) < 0,$$

and the boundary equilibrium point A is the unique and globally stable equilibrium of (4) in \mathbb{R}_+^3 .

From Lemma 2.1 we know that for any choice of initial conditions, the trajectory of system (4) always approaches to A when $R_0 < 1$. In other words, cancer will be controlled and tumor cells will be annihilated eventually as long as the increasing rate of lymphocyte cells is large enough.

Point B exists as a nonnegative equilibrium only if $R_0 \geq 1$. In this case, point A turns to be a saddle with a one-dimensional unstable manifold from a stable node, which implies that for each orbit starting in \mathbb{R}_p^3 the number of cancer cells will not tend to zero. A bifurcation occurs at $R_0 = 1$.

Remark 1. The role of R_0 is similar to the basic reproduction number in epidemic models.

Now we intend to study the qualitative properties of (4) near the equilibrium B . The Jacobian matrix of (4) at B is

$$J_B = \begin{pmatrix} -\lambda_1 & \lambda_1 x_B [\frac{1}{y_B(1+y_B)} - \frac{1}{y_c}] & \frac{\alpha_1 y_B}{1+y_B} \exp(-\frac{y_B}{y_c}) \\ \lambda_1 & -\alpha_3 & 0 \\ 0 & -\frac{\alpha_2 \lambda_1 x_B}{\alpha_1 y_B} \exp(\frac{y_B}{y_c}) & 0 \end{pmatrix} \tag{6}$$

and the corresponding characteristic equation is

$$\lambda^3 + a_1 \lambda^2 + a_2 \lambda + a_3 = 0, \tag{7}$$

where

$$\begin{aligned} a_1 &= \alpha_3 + \lambda_1 > 0, \\ a_2 &= \lambda_1^2 [\frac{x_0}{y_B} + x_B (\frac{1}{y_c} + \frac{1}{1+y_B})] > 0, \\ a_3 &= \frac{\alpha_2 \lambda_1^2 x_B}{1+y_B} > 0. \end{aligned}$$

Applying Routh-Hurwitz criterion to the Jacobian matrix J_B , we achieve that the sufficient condition for the stability of B is

$$a_1 a_2 - a_3 = \lambda_1^2 \{ (\alpha_3 + \lambda_1) [\frac{x_0}{y_B} + x_B (\frac{1}{y_c} + \frac{1}{1+y_B})] - \frac{\alpha_2 x_B}{1+y_B} \} > 0.$$

Thus, point B is unstable if $a_1 a_2 - a_3 < 0$ but is uncertain if $a_1 a_2 - a_3 = 0$. Nevertheless, equation (7) always has at least one negative real root γ no matter what the sign of $a_1 a_2 - a_3$ is. Also γ is the principle eigenvalue, and the absolute value of γ is greater than that of the real parts of the other eigenvalues.

Under the assumption $a_1 a_2 - a_3 = 0$, it is clear to see that J_B has one negative eigenvalue $-(\alpha_3 + \lambda_1)$ and two pure imaginary eigenvalues $\pm iw$, where $w > 0$ satisfies $w^2 = a_2 = \lambda_1^2 [\frac{x_0}{y_B} + x_B (\frac{1}{y_c} + \frac{1}{1+y_B})]$. Although the stability of B is not determined, Hopf bifurcation may occur at this equilibrium. We resort to the following Hopf bifurcation theorem in three-dimensional differential systems to determine the occurrence of Hopf bifurcation which can be found in Zhang [32].

Lemma 2.2. *Let $\Omega \subseteq R^3$ be an open set containing $O(x_1, y_1, z_1)$ and let $S \subseteq R$ be an open set with $0 \in S$. Let $f : \Omega \times S \rightarrow R^3$ be an analytic function such that $f(O, \mu) = 0$ for any $\mu \in S$. Assume the variational matrix $Df(O, \mu)$ of f has one real eigenvalue $\gamma(\mu)$ and two conjugate imaginary eigenvalues $\alpha(\mu) \pm i\beta(\mu)$ with $\gamma(0) < 0$, $\alpha(0) = 0$, $\beta(0) > 0$. Furthermore, suppose that the eigenvalues cross the imaginary axis with nonzero speed, that is, $\frac{d\alpha}{d\mu}(0) \neq 0$. Then the following differential system*

$$\dot{X} = f(X, \mu) \tag{8}$$

undergoes Hopf bifurcation near the equilibrium point O at $\mu = 0$.

Remark 2. If $\alpha'(0) > 0$ and O is a stable but not asymptotically stable equilibrium when $\mu = 0$, then the solutions of system (8) on some surface in the neighborhood of O are all periodic orbits.

Remark 3. If $\alpha'(0) > 0$ and O is a asymptotically stable (unstable) equilibrium when $\mu = 0$, then system (8) has an asymptotically stable periodic orbit in the neighborhood of O for sufficiently small $\mu > 0$ ($\mu < 0$).

In order to clarify how system (4) undergoes Hopf bifurcation at point B , we choose the death rate α_3 of stage-2 lymphocytes as a perturbed parameter. The equation $a_1 a_2 - a_3 = 0$ brings out

$$\alpha_3 = \frac{\alpha_2 y_B - \lambda_1(1 + y_B) \pm \sqrt{[\alpha_2 y_B - \lambda_1(1 + y_B)]^2 - 4\lambda_1 \alpha_2 x_0(1 + y_B)}}{2(1 + y_B)},$$

where we need $0 < \lambda_1 < \alpha_2 y_B / (1 + y_B)$ to guarantee that α_3 is nonnegative.

Without loss of generality, set $\alpha_3(\mu) = \alpha_3^0 + \mu$, where $\alpha_3(0) = \alpha_3$ satisfies that

$$(a_1 a_2 - a_3)|_{\mu=0} = \lambda_1^2 \left\{ (\alpha_3 + \lambda_1) \left[\frac{x_0}{y_B} + x_B \left(\frac{1}{y_c} + \frac{1}{1 + y_B} \right) \right] - \frac{\alpha_2 x_B}{1 + y_B} \right\} = 0. \quad (9)$$

We also need to determine the sign of the real part of $d\lambda/d\mu$ at $\mu = 0$ when the above equation is valid. Differentiating both sides of (7) with respect to μ , we have

$$\begin{aligned} & 3\lambda^2 \frac{d\lambda}{d\mu} + \lambda^2 + 2(\alpha_3 + \lambda_1) \lambda \frac{d\lambda}{d\mu} + \lambda_1^2 \left(\frac{1}{y_c} + \frac{1}{1 + y_B} \right) \lambda \frac{d x_B}{d\mu} \\ & + \lambda_1^2 \left[x_B \left(\frac{1}{y_c} + \frac{1}{1 + y_B} \right) + \frac{x_0}{y_B} \right] \frac{d\lambda}{d\mu} + \frac{\alpha_2 \lambda_1^2}{1 + y_B} \frac{d x_B}{d\mu} = 0, \end{aligned}$$

which leads to

$$\frac{d\lambda}{d\mu} = - \frac{\lambda^2 + \lambda_1 y_B \left(\frac{1}{y_c} + \frac{1}{1 + y_B} \right) \lambda + \frac{\alpha_2 \lambda_1 y_B}{1 + y_B}}{3\lambda^2 + 2(\alpha_3 + \lambda_1) \lambda + \lambda_1^2 \left[x_B \left(\frac{1}{y_c} + \frac{1}{1 + y_B} \right) + \frac{x_0}{y_B} \right]}, \quad (10)$$

where μ in x_B is omitted for simplicity. Therefore, by (9), we have

$$\begin{aligned} & \text{sign} \left\{ \text{Re} \left(\frac{d\lambda}{d\mu} \right) \Big|_{\mu=0} \right\} \\ & = \text{sign} \left\{ \text{Re} \left(- \frac{-w^2 + \lambda_1 y_B \left(\frac{1}{y_c} + \frac{1}{1 + y_B} \right) i w + \frac{\alpha_2 \lambda_1 y_B}{1 + y_B}}{-3w^2 + 2(\alpha_3 + \lambda_1) i w + \lambda_1^2 \left[x_B \left(\frac{1}{y_c} + \frac{1}{1 + y_B} \right) + \frac{x_0}{y_B} \right]} \right) \right\} \\ & = \text{sign} \left\{ - \left(-w^2 + \frac{\alpha_2 \lambda_1 y_B}{1 + y_B} \right) \left\{ -3w^2 + \lambda_1^2 \left[x_B \left(\frac{1}{y_c} + \frac{1}{1 + y_B} \right) + \frac{x_0}{y_B} \right] \right\} \right. \\ & \quad \left. - 2\lambda_1 (\alpha_3 + \lambda_1) y_B \left(\frac{1}{y_c} + \frac{1}{1 + y_B} \right) w^2 \right\} \\ & = \text{sign} \left\{ -w^2 + \frac{\alpha_2 \lambda_1 y_B}{1 + y_B} - \lambda_1 (\alpha_3 + \lambda_1) \left(\frac{1}{y_c} + \frac{1}{1 + y_B} \right) y_B \right\} \\ & = \text{sign} \left\{ - \frac{\lambda_1 \alpha_2 x_B}{(\alpha_3 + \lambda_1)(1 + y_B)} + \frac{x_0 (\alpha_3 + \lambda_1)}{x_B} \right\} \\ & = \text{sign} \left\{ -\lambda_1 \alpha_2 x_0^2 + [2\lambda_1 \alpha_2 \bar{x} + (\alpha_3 + \lambda_1)^2 (1 + y_B)] x_0 - \lambda_1 \alpha_2 \bar{x}^2 \right\} \\ & \triangleq \text{sign} \{ h(x_0) \}, \end{aligned} \quad (11)$$

where $\bar{x} = \frac{\alpha_3 \lambda_2}{\alpha_2 \lambda_1}$. Obviously, the degeneracy $\text{Re} \left(\frac{d\lambda}{d\mu} \right) \Big|_{\mu=0} = 0$ occurs at

$$x_0^* = \bar{x} + \frac{(\alpha_3 + \lambda_1)^2 (1 + y_B) - \sqrt{[2\lambda_1 \alpha_2 \bar{x} + (\alpha_3 + \lambda_1)^2 (1 + y_B)]^2 - 4\lambda_1^2 \alpha_2^2 \bar{x}^2}}{2\lambda_1 \alpha_2} < \bar{x},$$

when $h(x_0)$ vanishes (see Figure 1).




Figure 1. Graph of the function $h(x_0)$ defined on $[0, \bar{x}]$,
 $x_0^c = \bar{x} + \frac{(\alpha_3 + \lambda_1)^2(1 + y_B)}{2\lambda_1\alpha_2}$.

According to Lemma 2.2 and the above calculations, we obtain the following conclusion.

Theorem 2.3. *For any given $x_0 \in [0, \bar{x}]$ and $x_0 \neq x_0^*$, system (4) undergoes nondegenerate Hopf bifurcation at the nonhyperbolic equilibrium B when the parameters satisfy equation (9).*

When $x_0 = x_0^*$, the real part of $\frac{d\lambda}{d\mu}$ becomes zero, then the transversality condition fails. The occurrence of degeneracy may induce degenerate Hopf bifurcation or no Hopf bifurcation. We will continue to differentiate (10) associating to μ to determine the existence of degenerate Hopf bifurcation. Using (10), it can be easily deduced after tedious and mechanical calculations that

$$\begin{aligned} \frac{d^2\lambda}{d\mu^2} &= -\frac{[2\lambda + \lambda_1 y_B (\frac{1}{y_c} + \frac{1}{1+y_B})] \{3\lambda_1^2 + 2\lambda(\alpha_3 + \lambda_1) + \lambda_1^2 [x_B (\frac{1}{y_c} + \frac{1}{1+y_B}) + \frac{x_0}{y_B}]\} \frac{d\lambda}{d\mu}}{\{3\lambda^2 + 2\lambda(\alpha_3 + \lambda_1) + \lambda_1^2 [x_B (\frac{1}{y_c} + \frac{1}{1+y_B}) + \frac{x_0}{y_B}]\}^2} \\ &+ \frac{[\lambda^2 + \lambda_1 y_B (\frac{1}{y_c} + \frac{1}{1+y_B})\lambda + \frac{\alpha_2 \lambda_1 y_B}{1+y_B}] [6\lambda \frac{d\lambda}{d\mu} + 2(\alpha_3 + \lambda_1) \frac{d\lambda}{d\mu} + 2\lambda + \lambda_1 y_B (\frac{1}{y_c} + \frac{1}{1+y_B})]}{\{3\lambda^2 + 2\lambda(\alpha_3 + \lambda_1) + \lambda_1^2 [x_B (\frac{1}{y_c} + \frac{1}{1+y_B}) + \frac{x_0}{y_B}]\}^2}. \end{aligned} \tag{12}$$

On the assumption of (9) and the condition $x_0 = x_0^*$, it follows that $\text{Re}(\frac{d\lambda}{d\mu}|_{\mu=0}) = 0$ and

$$\begin{aligned} &\text{sign}\{\text{Re}(\frac{d^2\lambda}{d\mu^2})|_{\mu=0}\} \\ &= \text{sign}\{\text{Re}(\frac{[-w^2 + i\lambda_1 y_B (\frac{1}{y_c} + \frac{1}{1+y_B})]w + \frac{\alpha_2 \lambda_1 y_B}{1+y_B} [2iw + \lambda_1 y_B (\frac{1}{y_c} + \frac{1}{1+y_B})]}{\{-3w^2 + 2i(\alpha_3 + \lambda_1)w + \lambda_1^2 [x_B (\frac{1}{y_c} + \frac{1}{1+y_B}) + \frac{x_0}{y_B}]\}^2})\} \\ &= \text{sign}\{\text{Re}(\frac{(-w^2 + \frac{\alpha_2 \lambda_1 y_B}{1+y_B})\lambda_1 y_B (\frac{1}{y_c} + \frac{1}{1+y_B}) - 2\lambda_1 y_B (\frac{1}{y_c} + \frac{1}{1+y_B})w^2}{w^2 - (\alpha_3 + \lambda_1)^2 - 2i(\alpha_3 + \lambda_1)w} \\ &\quad + \frac{i[2(-w^2 + \frac{\alpha_2 \lambda_1 y_B}{1+y_B})w + \lambda_1^2 y_B^2 (\frac{1}{y_c} + \frac{1}{1+y_B})^2 w]}{w^2 - (\alpha_3 + \lambda_1)^2 - 2i(\alpha_3 + \lambda_1)w})\} \\ &= \text{sign}\{\text{Re}(\frac{(-3w^2 + \frac{\alpha_2 \lambda_1 y_B}{1+y_B})\lambda_1 y_B (\frac{1}{y_c} + \frac{1}{1+y_B}) + i[2(-w^2 + \frac{\alpha_2 \lambda_1 y_B}{1+y_B}) + \lambda_1^2 y_B^2 (\frac{1}{y_c} + \frac{1}{1+y_B})^2]w}{w^2 - (\alpha_3 + \lambda_1)^2 - 2i(\alpha_3 + \lambda_1)w})\} \\ &= \text{sign}\{\text{Re}(\frac{-3w^2 + \frac{\alpha_2 \lambda_1 y_B}{1+y_B} + i[2(\alpha_3 + \lambda_1) + \lambda_1 y_B (\frac{1}{y_c} + \frac{1}{1+y_B})]w}{w^2 - (\alpha_3 + \lambda_1)^2 - 2i(\alpha_3 + \lambda_1)w})\} \\ &= \text{sign}\{(-3w^2 + \frac{\alpha_2 \lambda_1 y_B}{1+y_B})[w^2 - (\alpha_3 + \lambda_1)^2] - 2(\alpha_3 + \lambda_1)[2(\alpha_3 + \lambda_1) \\ &\quad + \lambda_1 y_B (\frac{1}{y_c} + \frac{1}{1+y_B})]w^2\} \\ &= \text{sign}\{-w^4 - w^2(\alpha_3 + \lambda_1)^2 - \frac{[3w^2 + (\alpha_3 + \lambda_1)^2]\alpha_2 \lambda_1 y_B}{1+y_B}\} \\ &< 0, \end{aligned} \tag{13}$$

where the last equation comes from $h(x_0) = 0$ in (11), which implies that $\lambda_1 y_B (\alpha_3 + \lambda_1) (\frac{1}{y_c} + \frac{1}{1+y_B}) = -w^2 + \frac{\alpha_2 \lambda_1 y_B}{1+y_B}$.




Figure 2. Three cases of $\alpha(\mu)$ when $|\mu| > 0$ is small enough.

Theorem 2.4. *For parameters satisfying equation (9), there is no Hopf bifurcation for system (4) at the nonhyperbolic equilibrium B when $x_0 = x_0^*$.*

From Figure 2 we can see that if $x_0 = x_0^*$, then when μ passes through the origin from any direction, the interior equilibrium is always locally stable. In terms of the notations in Golubitsky and Langford [12], $\alpha'(0) = 0$ is equivalent to $a_\mu(0, 0) = 0$. Since at the interior equilibrium of model (4) $a_\mu(0, 0) = 0$ holds but $\alpha''(0) < 0$, the normal form of $a(x^2, \mu)$ corresponds to the cases (2) or (7) or (10) in Theorem 3.19 of [12] for codimension less than 3. However, by the stability analysis, the corresponding bifurcation diagram should be displayed as in Fig. 4.2 of that paper.

3. Normalized form of the Jacobian matrix. In this section we provide an applicable transformation to normalize the Jacobian matrix. Consider the following C^∞ system in $R_+^3 \times S$:

$$\begin{cases} \dot{x}_1 = f_1(x_1, x_2, x_3, \mu), \\ \dot{x}_2 = f_2(x_1, x_2, x_3, \mu), \\ \dot{x}_3 = f_3(x_1, x_2, x_3, \mu), \end{cases} \tag{14}$$

where $R_+^3 = \{(x_1, x_2, x_3)^T | x_i \geq 0, i = 1, 2, 3\}$ and $S \subseteq R$ is an open set containing 0. Assume system (14) has an isolated equilibrium $X^*(x_1^*, x_2^*, x_3^*)$ for any $\mu \in S$ and the variational matrix of $f = (f_1, f_2, f_3)^T$ at point X^* is denoted as $A = (a_{ij})_{3 \times 3}$. Assume A has one negative real eigenvalue γ and a pair of purely imaginary eigenvalues $\pm iw$ ($w > 0$) as $\mu = 0$. Then the following lemma gives the normalized form of A .

Lemma 3.1. ([4, pp.106]) *Let A be a real $n \times n$ matrix. Then there exists a real nonsingular matrix P such that $\tilde{A} = P^{-1}AP$ has the real canonical form consisting of real square matrices $A_1, \dots, A_k, B_1, \dots, B_m$ down the main diagonal. Each A_j has the form*

$$A_j = \begin{pmatrix} S_j & 0_2 & \cdots & 0_2 & 0_2 \\ I_2 & S_j & \cdots & 0_2 & 0_2 \\ 0_2 & I_2 & \cdots & 0_2 & 0_2 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0_2 & 0_2 & \cdots & I_2 & S_j \end{pmatrix}, \tag{15}$$

where 0_2 is the 2×2 zero matrix, I_2 is the 2×2 identity matrix, and

$$S_j = \begin{pmatrix} \alpha_j & -\beta_j \\ \beta_j & \alpha_j \end{pmatrix}.$$

Then B_j has the form

$$B_j = \begin{pmatrix} \lambda_j & 0 & \cdots & 0 & 0 \\ 1 & \lambda_j & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & \lambda_j \end{pmatrix}.$$

We can validate this lemma by considering the canonical form of the above-mentioned 3×3 matrix A for $\mu = 0$ in explicit transformations. In fact, set

$$B(\lambda) \triangleq \lambda I_3 - A = \begin{pmatrix} \lambda - a_{11} & -a_{12} & -a_{13} \\ -a_{21} & \lambda - a_{22} & -a_{23} \\ -a_{31} & -a_{32} & \lambda - a_{33} \end{pmatrix},$$

where I_i is the $i \times i$ identity matrix. Then obviously the determinant of B is $\det B(\lambda) = (\lambda - \rho)(\lambda^2 + w^2)$. Moreover, we have

$$\begin{aligned} \rho &= a_{11} + a_{22} + a_{33}, \\ \rho w^2 &= \det A, \\ w^2 &= a_{11}a_{22} + a_{22}a_{33} + a_{11}a_{33} - a_{12}a_{21} - a_{23}a_{32} - a_{13}a_{31}. \end{aligned} \tag{16}$$

Assume $\beta = (\beta_1, \beta_2, \beta_3)^T$ is the right eigenvector of A associated to the eigenvalue ρ , then $A\beta = \rho\beta$, which is equivalent to the following equations:

$$\begin{cases} (\rho - a_{11})\beta_1 - a_{12}\beta_2 - a_{13}\beta_3 = 0, \\ -a_{21}\beta_1 + (\rho - a_{22})\beta_2 - a_{23}\beta_3 = 0, \\ -a_{31}\beta_1 - a_{32}\beta_2 + (\rho - a_{33})\beta_3 = 0. \end{cases} \tag{17}$$

Since the dimension of the basis of the eigenvector space corresponding to ρ is one, $\text{Rank}(B(\rho)) = 2$. Without loss of generality, let us assume

$$\det \begin{pmatrix} \rho - a_{11} & -a_{12} \\ -a_{21} & \rho - a_{22} \end{pmatrix} \neq 0.$$

Consider the first two equations of (17), we can solve one of the nonzero solutions as

$$(\beta_1, \beta_2, \beta_3)^T = \left(\left| \begin{array}{cc|c} a_{13} & -a_{12} & \rho - a_{11} \\ a_{23} & \rho - a_{22} & -a_{21} \end{array} \right|, \left| \begin{array}{cc|c} \rho - a_{11} & a_{13} & -a_{21} \\ -a_{21} & a_{23} & \rho - a_{22} \end{array} \right|, \left| \begin{array}{cc|c} \rho - a_{11} & -a_{12} & -a_{21} \\ -a_{21} & \rho - a_{22} & a_{23} \end{array} \right| \right)^T.$$

The following choice of a series of linear transformations is motivated by the methods in Lu and Luo [23]. Choose

$$T_1 = \begin{pmatrix} \rho - a_{11} & -a_{12} & \beta_1 \\ -a_{12} & \rho - a_{22} & \beta_2 \\ -a_{13} & -a_{23} & \beta_3 \end{pmatrix},$$

then we obtain that

$$T_1^{-1}AT_1 = \begin{pmatrix} c_{11} & c_{12} & 0 \\ c_{21} & c_{22} & 0 \\ c_{31} & c_{32} & \rho \end{pmatrix} \triangleq \begin{pmatrix} C_1 & 0 \\ C_2 & \rho \end{pmatrix},$$

where the elements of C_1 satisfy $c_{11} + c_{22} = 0$ and $w^2 = c_{11}c_{22} - c_{12}c_{21} > 0$.

Choose

$$T_2 = \begin{pmatrix} I_2 & 0 \\ C_2(C_1 - \rho I_2)^{-1} & 1 \end{pmatrix},$$

then

$$T_2^{-1}T_1^{-1}AT_1T_2 = \begin{pmatrix} C_1 & 0 \\ 0 & \rho \end{pmatrix}.$$

Take

$$T_3 = \begin{pmatrix} 1 & -c_{11}/w & 0 \\ 0 & -c_{21}/w & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

and set $T = T_1T_2T_3$, then we obtain

$$T^{-1}AT = \begin{pmatrix} o & w & 0 \\ -w & 0 & 0 \\ 0 & 0 & \rho \end{pmatrix}. \quad (18)$$

The lemma is proved.

4. Stability of periodic orbits. In order to determine the stability of periodic orbits bifurcated from the interior equilibrium of system (4), we apply the general formula in Hsü and Kazarinoff [14], which turns out to be more applicable than the original bifurcation formulas given by Hopf [13] and Friedrichs [11], and the center manifold theorem in Perko [24]. Firstly we need to outline this method and some notations. Consider the following class of autonomous differential systems:

$$\dot{X} = A(\mu)X + F(X, \mu), \quad (19)$$

where $X = (x_1, x_2, \dots, x_n)^T$, F is a real analytic function on $\Omega \times (-c, c)$ with $F(0, \mu) \equiv 0$ and $F_X(0, \mu) \equiv 0$, Ω is an open connected domain in \mathbb{R}^n , $c > 0$ and A is a real $n \times n$ analytic matrix defined on $(-c, c)$ with exactly two purely imaginary eigenvalues at $\mu = 0$ whose continuous extensions are denoted by $\alpha(\mu)$ and $\bar{\alpha}(\mu)$ such that

$$\alpha(0) = -\bar{\alpha}(0), \quad \operatorname{Re}(\alpha'(0)) \neq 0, \quad \operatorname{Im}(\alpha(0)) = w > 0. \quad (20)$$

In virtue of Lemma 3.1, there exists a nonsingular matrix P which can transform A into the normal form with the Jordan block as A_j and B_j . For the sake of simplicity, in the following discussion we suppose A has the normal form as

$$A(0) = \begin{pmatrix} 0 & w & 0 \\ -w & 0 & 0 \\ 0 & 0 & D \end{pmatrix}. \quad (21)$$

Hopf [13] obtained the bifurcation theorem on the existence of a periodic analytic solution $p(t, \epsilon)$ with period $T(\epsilon)$ of system (26) under the above assumptions, where the parameter ϵ is related to μ by an analytic function $\mu = \mu(\epsilon)$ such that $\mu(0) = 0$, $p(t, 0) = 0$, $T(0) = 2\pi/w$ and $p(t, \epsilon) \neq 0$ for all sufficiently small $|\epsilon| > 0$. Furthermore, these periodic solutions exist for exactly one of three cases: $\mu > 0$, $\mu < 0$, or $\mu = 0$. By Hopf's theorem, the characteristic exponents of this bifurcated periodic solution $p(t, \epsilon)$ are the eigenvalues of the eigenvalue problem

$$\dot{V}(t) + \lambda V(t) = L(t, \epsilon)V(t), \quad (22)$$

where $V(t)$ has the same period $T(\epsilon)$ and $L(t, \epsilon) = A(\mu(\epsilon)) + F_X(p(t, \epsilon), \mu(\epsilon))$. According to the assumptions, there are exactly two characteristic exponents, which depend continuously upon ϵ , are determined only by $\operatorname{mod}(2\pi i/T(\epsilon))$ and tend to the imaginary axis as $\epsilon \rightarrow 0$, one of which is identically zero and the other $\beta = \beta(\epsilon)$

must be real and analytic at $\epsilon = 0$ satisfying $\beta(0) = 0$. Hopf [13] pointed out that in the expansions

$$\mu = \mu_1\epsilon + \mu_2\epsilon^2 + \dots, \quad \beta = \beta_1\epsilon + \beta_2\epsilon^2 + \dots, \tag{23}$$

the coefficients μ_1 and β_1 must be both zeros and there are the elementary relationships between μ_2 and β_2 :

$$\begin{aligned} \beta_2 &= -2\mu_2\text{Re}(\alpha'(0)) \\ &= 2 \int_0^{T_0} [F_{XX}(X^0, X^1) + \frac{1}{6}F_{XXX}(X^0, X^0, X^0)] \cdot Z(t)dt, \end{aligned} \tag{24}$$

where X^0, X^1 are T_0 -periodic solutions of systems

$$\dot{X}^0 = A(0)X^0 \tag{25}$$

and

$$\dot{X}^1 = A(0)X^1 + \frac{1}{2}F_{XX}(X^0, X^0), \tag{26}$$

respectively, here

$$F_{XX}(X^0, X^1) = \sum_{i,j}^n \frac{\partial^2 F(0,0)}{\partial x_i \partial x_j} x_i^0 x_j^1 \triangleq \sum_{i,j}^n F_{ij} x_i^0 x_j^1, \tag{27}$$

$$F_{XXX}(X^0, X^0) = \sum_{i,j}^n \frac{\partial^3 F(0,0)}{\partial x_i \partial x_j} x_i^0 x_j^0 \triangleq \sum_{i,j}^n F_{ij} x_i^0 x_j^0, \tag{28}$$

$Z(t)$ is a T_0 -periodic solution of

$$\dot{Z}(t) = -A^T(0)Z(t), \tag{29}$$

and $T_0 = T(0)$ such that

$$\int_0^{T_0} X^0(s)Z(s)ds = 1 \quad \text{and} \quad \int_0^{T_0} \dot{X}^0(s)Z(s)ds = 0.$$

The interested readers are referred to Hopf [13] for more details.

Based on the assumptions on A , $A(0)$ has a pair of simple complex conjugate eigenvalues $\pm wi$, thus any T_0 -periodic solution of (25) is a linear combination of $e^{wit}a(0)$ and $e^{wit}a(0)$, where $a(0) = (1, i, 0, \dots, 0)^T$ is the characteristic vector corresponding to $\alpha(0)$. For the autonomous system (25), any nontrivial T_0 -periodic solution $X^0(t)$ may be assumed as the form $X^0(t) = (b \cos(\omega t), -b \sin(\omega t), 0, \dots, 0)^T$ ($b > 0$) (see Poore [25, (4.7)]). For the nonlinear autonomous system (26), Hsü and Kazarinoff [14] proved that there exists a unique T_0 -periodic solution $X^1(t)$ corresponding to X^0 , which can be written as

$$X^1 = \frac{b^2}{6w}(\cos(\omega t)h_1 + \sin(\omega t)h_2, -\sin(\omega t)h_1 + \cos(\omega t)h_2, (\tilde{X}^1(t))^T)^T, \tag{30}$$

where

$$\begin{aligned} h_1 &= \sin^3(\omega t)(-F_{11}^1 + 2F_{12}^2 + F_{22}^1) + \cos^3(\omega t)(F_{11}^1 + 2F_{12}^1 - F_{22}^2) \\ &\quad + 3[\sin(\omega t)F_{11}^1 + \cos(\omega t)F_{22}^2] + 2F_{11}^2 - 2F_{12}^1 - F_{22}^2, \\ h_2 &= -\sin^3(\omega t)(F_{11}^2 + 2F_{12}^1 - F_{22}^2) + \cos^3(\omega t)(-F_{11}^1 + 2F_{12}^2 + F_{22}^1) \\ &\quad + 3[\sin(\omega t)F_{11}^2 - \cos(\omega t)F_{22}^1] + F_{11}^1 - 2F_{12}^2 + 2F_{22}^1, \\ \tilde{X}^1(t) &= -\frac{3w}{2}D^{-1}(\tilde{F}_{11} + \tilde{F}_{22}) \\ &\quad + \frac{3}{8w}(I + \frac{D^2}{4w^2})^{-1}[2w \sin(2\omega t)I - \cos(2\omega t)D](\tilde{F}_{11} - \tilde{F}_{22}) \\ &\quad + \frac{3}{4w}(I + \frac{D^2}{4w^2})^{-1}[2w \cos(2\omega t)I + \sin(2\omega t)D]\tilde{F}_{12}, \end{aligned}$$

where $\tilde{X}^1(t) = (x_3, x_4, \dots, x_n)^T$, $\tilde{F}_{ij} = (\frac{\partial^2 F^3(0,0)}{\partial x_i \partial x_j}, \frac{\partial^2 F^4(0,0)}{\partial x_i \partial x_j}, \dots, \frac{\partial^2 F^n(0,0)}{\partial x_i \partial x_j})^T$ and F_{ij}^k is the k th component of F_{ij} ($i, j, k = 1, 2$), D is defined as in (21) and I is the $(n - 2) \times (n - 2)$ identity matrix.

Based on the expressions (24), (30) and an algebraic computation, Hsü and Kazarinoff [14] derived a formula for the direction of the bifurcation of periodic orbits of system (19).

Lemma 4.1. ([14]) *If $\mu \neq 0$, the direction of the bifurcation of (4) is determined by the following equation,*

$$\begin{aligned} \mu_2 = & \frac{b_0^2}{16\text{Re}(\alpha'(0))} \left\{ \frac{1}{w} [(F_{11}^1 + F_{22}^1)F_{12}^1 - (F_{11}^2 + F_{22}^2)F_{12}^2 - F_{11}^1 F_{11}^2 + F_{22}^1 F_{22}^2] \right. \\ & - (F_{111}^1 + F_{122}^1 + F_{112}^2 + F_{222}^2) + 2(G_1^1 + G_2^2)D^{-1}(\bar{F}_{11} + \bar{F}_{22}) \\ & + \frac{1}{2w^2}(G_2^1 + G_1^2)\left(I + \frac{D^2}{4w^2}\right)^{-1}[w(\bar{F}_{11} - \bar{F}_{22}) + D\bar{F}_{12}] \\ & \left. + \frac{1}{4w^2}(G_1^1 - G_2^2)\left(I + \frac{D^2}{4w^2}\right)^{-1}[D(\bar{F}_{11} - \bar{F}_{22}) - 4w\bar{F}_{12}] \right\}, \end{aligned} \tag{31}$$

where $G_j^i = (F_{j,3}^i, F_{j,4}^i, \dots, F_{j,n}^i)$ for $i,j=1,2$, and $(b, 0, \dots, 0)^T$ is the initial value of X_0 .

The stability of the periodic orbit bifurcated from the degenerate equilibrium can be determined by the following lemma.

Lemma 4.2. ([14]) *Under the assumptions in Hopf’s theorem, if $A(0)$ has exactly two purely imaginary eigenvalues and the other $n - 2$ eigenvalues have negative real part and if $\mu_2\text{Re}(\alpha'(0)) > 0$, then a bifurcating periodic solution whose existence is asserted by Hopf’s theorem is asymptotically orbitally stable with asymptotic phase; however, if $\mu_2\text{Re}(\alpha'(0)) < 0$, these small periodic solutions are unstable. Moreover, if any one of the other $n - 2$ eigenvalues has positive real part, then the bifurcating periodic solution is orbitally unstable.*

Now let us return to determine the stability of limit cycles bifurcated from the equilibrium B of system (4). It is necessary first to transform the Jacobian matrix J_B expressed in (6) into the normal form. For simplicity, let us set

$$a \triangleq \lambda_1 x_B \left[\frac{1}{y_B(1+y_B)} - \frac{1}{y_c} \right], \quad b \triangleq \frac{\alpha_1 y_B}{1+y_B} \exp\left(-\frac{y_B}{y_c}\right), \quad c \triangleq \frac{\alpha_2 \lambda_1 x_B}{\alpha_1 y_B} \exp\left(\frac{y_B}{y_c}\right) = \frac{\alpha_2 z_B}{1+y_B}, \tag{32}$$

where b and c are both positive, then J_B can be written as

$$J_B \triangleq \begin{pmatrix} -\lambda_1 & a & b \\ \lambda_1 & -\alpha_3 & 0 \\ 0 & -c & 0 \end{pmatrix}.$$

Suppose system (4) undergoes Hopf bifurcation at point B , then the parameters in J_B satisfy

$$bc = (\alpha_3 - a)(\lambda_1 + \alpha_3) \tag{33}$$

and J_B has one negative eigenvalue $-\delta = -(\lambda_1 + \alpha_3)$ and a pair of purely imaginary eigenvalues $\pm wi$, where $w^2 = \lambda_1(\alpha_3 - a) > 0$. Select

$$P = \begin{pmatrix} \alpha_3 & w & -1 \\ \lambda_1 & 0 & 1 \\ 0 & c\lambda_1 w^{-1} & (\alpha_3 - a)b^{-1} \end{pmatrix}.$$

It is easy to check that the inverse matrix of P turns out to be

$$P^{-1} = \Delta^{-1} \begin{pmatrix} c\lambda_1 w^{-1} & w(\alpha_3 - a)b^{-1} + c\lambda_1 w^{-1} & -w \\ w^2 b^{-1} & -\alpha_3(\alpha_3 - a)b^{-1} & \delta \\ -c\lambda_1^2 w^{-1} & c\lambda_1 \alpha_3 w^{-1} & w\lambda_1 \end{pmatrix}$$

and $L(0) \triangleq P^{-1}J_B P$ becomes the normal form as the right side in (18), where $\Delta = w(w^2 + \delta^2)/b$.

In the following a series of linear transformations will be introduced for system (4). Let $x_1 = x - x_B$, $y_1 = y - y_B$, $z_1 = z - z_B$, then the positive equilibrium B is translated into the origin of the system:

$$\begin{aligned}
 & (\dot{x}_1, \dot{y}_1, \dot{z}_1)^T \\
 = & J_B \begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} + \begin{pmatrix} \frac{1}{2}f_{yy}y_1^2 + f_{yz}y_1z_1 + \frac{1}{3!}f_{yyy}y_1^3 + \frac{1}{2}f_{yyz}y_1^2z_1 + Q_1^1(y_1, z_1) \\ 0 \\ \frac{1}{2}h_{yy}y_1^2 + h_{yz}y_1z_1 + \frac{1}{3!}h_{yyy}y_1^3 + \frac{1}{2}h_{yyz}y_1^2z_1 + Q_1^3(y_1, z_1) \end{pmatrix},
 \end{aligned} \tag{34}$$

where $Q_1^i (i = 1, 3)$ are C^∞ power series of y_1 and z_1 with power higher than 3, f_{yy} , f_{yz} , f_{yyy} , f_{yyz} and h_{yy} , h_{yz} , h_{yyy} , h_{yyz} are the corresponding partial derivatives at point B .

If we denote $X_i = (x_i, y_i, z_i)^T (i = 1, 2)$ and let $X_2 = P^{-1}X_1$, then

$$\begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} = \begin{pmatrix} \alpha_3x_2 + wy_2 - z_2 \\ \lambda_1x_2 + z_2 \\ c\lambda_1w^{-1}y_2 + (\alpha_3 - a)b^{-1}z_2 \end{pmatrix},$$

and system (34) is changed into

$$\begin{aligned}
 \dot{X}_2 &= P^{-1}\dot{X}_1 = P^{-1}J_BPX_2 \\
 &+ P^{-1} \begin{pmatrix} \frac{1}{2}f_{yy}y_1^2 + f_{yz}y_1z_1 + \frac{1}{3!}f_{yyy}y_1^3 + \frac{1}{2}f_{yyz}y_1^2z_1 + Q_1^1(y_1, z_1) \\ 0 \\ \frac{1}{2}h_{yy}y_1^2 + h_{yz}y_1z_1 + \frac{1}{3!}h_{yyy}y_1^3 + \frac{1}{2}h_{yyz}y_1^2z_1 + Q_1^3(y_1, z_1) \end{pmatrix} \\
 &\triangleq L(0)X_2 + P^{-1}G(x_2, y_2, z_2),
 \end{aligned} \tag{35}$$

where

$$\begin{aligned}
 G &= \begin{pmatrix} \frac{1}{2}f_{yy}(\lambda_1x_2 + z_2)^2 + f_{yz}(\lambda_1x_2 + z_2)[c\lambda_1w^{-1}y_2 + (\alpha_3 - a)b^{-1}z_2] \\ + \frac{1}{2}f_{yyz}(\lambda_1x_2 + z_2)^2[c\lambda_1w^{-1}y_2 + (\alpha_3 - a)b^{-1}z_2] \\ + \frac{1}{3!}f_{yyy}(\lambda_1x_2 + z_2)^3 + Q_2^1(x_2, y_2, z_2) \\ 0 \\ \frac{1}{2}h_{yy}(\lambda_1x_2 + z_2)^2 + h_{yz}(\lambda_1x_2 + z_2)[c\lambda_1w^{-1}y_2 + (\alpha_3 - a)b^{-1}z_2] \\ + \frac{1}{2}h_{yyz}(\lambda_1x_2 + z_2)^2[c\lambda_1w^{-1}y_2 + (\alpha_3 - a)b^{-1}z_2] \\ + \frac{1}{3!}h_{yyy}(\lambda_1x_2 + z_2)^3 + Q_2^3(x_2, y_2, z_2) \end{pmatrix} \\
 &= \begin{pmatrix} \frac{\lambda_1^2}{2}f_{yy}x_2^2 + c\lambda_1^2w^{-1}f_{yz}x_2y_2 + \lambda_1[f_{yy} + (\alpha_3 - a)b^{-1}f_{yz}]x_2z_2 \\ + c\lambda_1w^{-1}f_{yz}y_2z_2 + [\frac{1}{2}f_{yy} + (\alpha_3 - a)b^{-1}f_{yz}]z_2^2 \\ + \frac{1}{3!}\lambda_1^3f_{yyy}x_2^3 + \frac{\lambda_1^2}{2}[f_{yyy} + (\alpha_3 - a)b^{-1}f_{yyz}]x_2^2z_2 \\ + \frac{c\lambda_1^3w^{-1}}{2}f_{yyz}x_2^2y_2 + \lambda_1[\frac{1}{2}f_{yyy} + (\alpha_3 - a)b^{-1}f_{yyz}]x_2z_2^2 \\ + \frac{c\lambda_1w^{-1}}{2}f_{yyz}y_2z_2^2 + [\frac{1}{3!}f_{yyy} + \frac{1}{2}(\alpha_3 - a)b^{-1}f_{yyz}]z_2^3 \\ + c\lambda_1^2w^{-1}f_{yyz}x_2y_2z_2 + Q_2^1(x_2, y_2, z_2) \\ 0 \\ \frac{\lambda_1^2}{2}h_{yy}x_2^2 + c\lambda_1^2w^{-1}h_{yz}x_2y_2 + \lambda_1[h_{yy} + (\alpha_3 - a)b^{-1}h_{yz}]x_2z_2 \\ + c\lambda_1w^{-1}h_{yz}y_2z_2 + [\frac{1}{2}h_{yy} + (\alpha_3 - a)b^{-1}h_{yz}]z_2^2 \\ + \frac{1}{3!}\lambda_1^3h_{yyy}x_2^3 + \frac{\lambda_1^2}{2}[h_{yyy} + (\alpha_3 - a)b^{-1}h_{yyz}]x_2^2z_2 \\ + \frac{c\lambda_1^3w^{-1}}{2}h_{yyz}x_2^2y_2 + \lambda_1[\frac{1}{2}h_{yyy} + (\alpha_3 - a)b^{-1}h_{yyz}]x_2z_2^2 \\ + \frac{c\lambda_1w^{-1}}{2}h_{yyz}y_2z_2^2 + [\frac{1}{3!}h_{yyy} + \frac{1}{2}(\alpha_3 - a)b^{-1}h_{yyz}]z_2^3 \\ + c\lambda_1^2w^{-1}h_{yyz}x_2y_2z_2 + Q_2^3(x_2, y_2, z_2) \end{pmatrix},
 \end{aligned}$$

in which $Q_2^i (i = 1, 3)$ are C^∞ power series of x_2, y_2, z_2 with power higher than 3.

Denote $G = (G^1, G^2, G^3)^T$ and $F = P^{-1}G$, then

$$F = \Delta^{-1} \begin{pmatrix} c\lambda_1 w^{-1} G^1 - wG^3 \\ w^2 b^{-1} G^1 + \delta G^3 \\ -c\lambda_1^2 w^{-1} G^1 + w\lambda_1 G^3 \end{pmatrix} \triangleq \begin{pmatrix} F^1 + Q_3^1(x_2, y_2, z_2) \\ F^2 + Q_3^2(x_2, y_2, z_2) \\ F^3 + Q_3^3(x_2, y_2, z_2) \end{pmatrix},$$

where

$$\begin{aligned} F^1 &= \Delta^{-1} \{ \frac{\lambda_2^2}{2} (c\lambda_1 w^{-1} f_{yy} - wh_{yy}) x_2^2 + c\lambda_1^2 w^{-1} (c\lambda_1 w^{-1} f_{yz} - wh_{yz}) x_2 y_2 \\ &\quad + \lambda_1 [c\lambda_1 w^{-1} f_{yy} - wh_{yy} + (\alpha_3 - a) b^{-1} (c\lambda_1 w^{-1} f_{yz} - wh_{yz})] x_2 z_2 \\ &\quad + c\lambda_1 w^{-1} (c\lambda_1 w^{-1} f_{yz} - wh_{yz}) y_2 z_2 \\ &\quad + [\frac{1}{2} (c\lambda_1 w^{-1} f_{yy} - wh_{yy}) + (\alpha_3 - a) b^{-1} (c\lambda_1 w^{-1} f_{yz} - wh_{yz})] z_2^2 \\ &\quad + \frac{1}{3!} \lambda_1^3 (c\lambda_1 w^{-1} f_{yyy} - wh_{yyy}) x_2^3 + \frac{c\lambda_1^3 w^{-1}}{2} (c\lambda_1 w^{-1} f_{yyz} - wh_{yyz}) x_2^2 y_2 \\ &\quad + \frac{\lambda_2^2}{2} [(c\lambda_1 w^{-1} f_{yyy} - wh_{yyy}) + (\alpha_3 - a) b^{-1} (c\lambda_1 w^{-1} f_{yyz} - wh_{yyz})] x_2^2 z_2 \\ &\quad + c\lambda_1^2 w^{-1} (c\lambda_1 w^{-1} f_{yyz} - wh_{yyz}) x_2 y_2 z_2 \\ &\quad + \lambda_1 [\frac{1}{2} (c\lambda_1 w^{-1} f_{yyy} - wh_{yyy}) + (\alpha_3 - a) b^{-1} (c\lambda_1 w^{-1} f_{yyz} - wh_{yyz})] x_2 z_2^2 \\ &\quad + \frac{c\lambda_1 w^{-1}}{2} (c\lambda_1 w^{-1} f_{yyz} - wh_{yyz}) y_2 z_2^2 \\ &\quad + [\frac{1}{3!} (c\lambda_1 w^{-1} f_{yyy} - wh_{yyy}) + \frac{1}{2} (\alpha_3 - a) b^{-1} (c\lambda_1 w^{-1} f_{yyz} - wh_{yyz})] z_2^3 \}, \\ F^2 &= \Delta^{-1} \{ \frac{\lambda_2^2}{2} (w^2 b^{-1} f_{yy} + \delta h_{yy}) x_2^2 + c\lambda_1^2 w^{-1} (w^2 b^{-1} f_{yz} + \delta h_{yz}) x_2 y_2 \\ &\quad + \lambda_1 [w^2 b^{-1} f_{yy} + \delta h_{yy} + (\alpha_3 - a) b^{-1} (w^2 b^{-1} f_{yz} + \delta h_{yz})] x_2 z_2 \\ &\quad + c\lambda_1 w^{-1} (w^2 b^{-1} f_{yz} + \delta h_{yz}) y_2 z_2 \\ &\quad + [\frac{1}{2} (w^2 b^{-1} f_{yy} + \delta h_{yy}) + (\alpha_3 - a) b^{-1} (w^2 b^{-1} f_{yz} + \delta h_{yz})] z_2^2 \\ &\quad + \frac{1}{3!} \lambda_1^3 (w^2 b^{-1} f_{yyy} + \delta h_{yyy}) x_2^3 + \frac{c\lambda_1^3 w^{-1}}{2} (w^2 b^{-1} f_{yyz} + \delta h_{yyz}) x_2^2 y_2 \\ &\quad + \frac{\lambda_2^2}{2} [(w^2 b^{-1} f_{yyy} + \delta h_{yyy}) + (\alpha_3 - a) b^{-1} (w^2 b^{-1} f_{yyz} + \delta h_{yyz})] x_2^2 z_2 \\ &\quad + c\lambda_1^2 w^{-1} (w^2 b^{-1} f_{yyz} + \delta h_{yyz}) x_2 y_2 z_2 \\ &\quad + \lambda_1 [\frac{1}{2} (w^2 b^{-1} f_{yyy} + \delta h_{yyy}) + (\alpha_3 - a) b^{-1} (w^2 b^{-1} f_{yyz} + \delta h_{yyz})] x_2 z_2^2 \\ &\quad + \frac{c\lambda_1 w^{-1}}{2} (w^2 b^{-1} f_{yyz} + \delta h_{yyz}) y_2 z_2^2 \\ &\quad + [\frac{1}{3!} (w^2 b^{-1} f_{yyy} + \delta h_{yyy}) + \frac{1}{2} (\alpha_3 - a) b^{-1} (w^2 b^{-1} f_{yyz} + \delta h_{yyz})] z_2^3 \}, \\ F^3 &= -\lambda_1 F^1, \end{aligned} \tag{36}$$

and $Q_3^1 = c\lambda_1 w^{-1} Q_2^1 - wQ_2^3$, $Q_3^2 = w^2 b^{-1} Q_2^1 + \delta Q_2^3$, $Q_3^3 = -\lambda_1 Q_3^1$ are C^∞ power series of x_2, y_2, z_2 with power higher than 3.

Let $F_1^i = \frac{\partial F^i}{\partial x_2}(0)$, $F_2^i = \frac{\partial F^i}{\partial y_2}(0)$, $F_3^i = \frac{\partial F^i}{\partial z_2}(0)$, then F_{22}^i , F_{122}^i , F_{222}^i are all vanished for $i = 1, 2, 3$. We obtain the following lemma directly from Lemma (4.1).

Lemma 4.3. *For system (35), it holds that*

$$\begin{aligned} \frac{16\mu_2 \text{Re}(\alpha')}{b_0^2} &= w^{-1} (F_{11}^1 F_{12}^1 - F_{11}^2 F_{12}^2 - F_{11}^1 F_{11}^2) - (F_{111}^1 + F_{112}^2) \\ &\quad - \frac{\delta}{4w^2 + \delta^2} [(F_{13}^1 - F_{23}^2) F_{11}^3 + 2(F_{23}^1 + F_{13}^2) F_{12}^3] - \frac{2}{\delta} (F_{13}^1 + F_{23}^2) F_{11}^3 \\ &\quad + \frac{2w}{4w^2 + \delta^2} [(F_{13}^1 + F_{13}^2) F_{11}^3 - 2(F_{13}^1 - F_{23}^2) F_{12}^3]. \end{aligned}$$

In order to get a stable periodic orbit bifurcated from the equilibrium B , we constrain our discussion on the hypothesis $y_c = y_B(1 + y_B)$ in this section. As a matter of fact, it is obvious to find from (36) the equations as below,

$$\begin{aligned} F_{13}^1 &= \frac{1}{\lambda_1} F_{11}^1 + \frac{(\alpha_3 - a)b^{-1}w}{c\lambda_1} F_{12}^1, \quad F_{23}^1 = \frac{1}{\lambda_1} F_{12}^1, \quad F_{13}^2 = \frac{1}{\lambda_1} F_{11}^2 - \frac{1}{\lambda_1} F_{12}^1, \\ F_{23}^2 &= -\frac{w^{-1}(\alpha_3 + \lambda_1)}{\lambda_1} F_{12}^1, \quad F_{11}^3 = -\lambda_1 F_{11}^1, \quad F_{12}^3 = -\lambda_1 F_{12}^1, \quad F_{112}^2 = \frac{c\lambda_1 w^{-1}}{z_B} F_{11}^2. \end{aligned}$$

Therefore, the next conclusion is followed from Lemma 4.3 after some simple calculations.

Lemma 4.4. *Assume $y_c = y_B(1 + y_B)$, then for system (35) it holds that*

$$\begin{aligned} \frac{16\mu_2\text{Re}(\alpha'(0))}{b_0^2} &= \left(\frac{\delta}{4w^2+\delta^2} + \frac{2}{\delta}\right)(F_{11}^1)^2 + \left[\frac{4w^2}{\delta(4w^2+\delta^2)} + \frac{4\delta}{4w^2+\delta^2}\right](F_{12}^1)^2 \\ &+ \frac{\delta(6w^2+\delta^2)}{w^2(4w^2+\delta^2)}F_{12}^1F_{11}^1 - \frac{c\lambda_1}{wz_B}F_{11}^2 - \frac{6w^2+\delta^2}{w(4w^2+\delta^2)}F_{11}^1F_{11}^2 \\ &+ \frac{w(8w^2+3\delta^2)}{\delta^2(4w^2+\delta^2)}F_{11}^1F_{12}^1 - F_{111}^1. \end{aligned} \tag{37}$$

Now let us set

$$\begin{aligned} H_1 &= y_B^2[\lambda_2^2(\delta^2\alpha_2^2 - \delta^4 - \delta^2w^2 - 4w^4) + \delta(\alpha_2 + 2\lambda_2)(\delta^4 + 7\delta^2w^2 + 2w^4)], \\ H_2 &= \delta^2w^4(1 + 2y_B)^2 + \delta\lambda_2(\delta^4 + 5\delta^2w^2 + 4w^4)(1 + 3y_B + 3y_B^2). \end{aligned}$$

Then the next theorem on the bifurcation of a stable periodic orbit will be achieved.

Theorem 4.5. *Assume that the parameters in (4) satisfy the degenerate condition (9) and $y_c = y_B(1 + y_B)$. If $\alpha_2^2 \leq 2\lambda_2^2$ and $H_1 > H_2$, then when α_3 undergoes slightly small perturbations, an asymptotically stable limit cycle will be bifurcated from the Hopf bifurcation point B in (4).*

Proof. From the expressions of f and h in (4), it can be easily found that $f_{yz} = 0$ in the case of $y_c = y_B(1 + y_B)$. We also obtain that

$$\begin{aligned} f_{yy} &= -\frac{\alpha_1z_B(1+2y_B)e^{-\frac{y_B}{y_c}}}{y_B(1+y_B)^3} = -\frac{\lambda_1(1+2y_B)x_B}{y_B^2(1+y_B)^2} < 0, \\ h_{yy} &= \frac{2(\alpha_2+\lambda_2)z_B}{(1+y_B)^3} > 0, \quad h_{yz} = -\frac{\alpha_2+\lambda_2}{(1+y_B)^2} < 0, \\ f_{yyy} &= \frac{\alpha_1z_B(2+6y_B+6y_B^2)e^{-\frac{y_B}{y_c}}}{y_B^3(1+y_B)^4} = -\frac{2+6y_B+6y_B^2}{y_B(1+y_B)(1+2y_B)}f_{yy} > 0, \\ f_{yyz} &= -\frac{\alpha_1(1+2y_B)e^{-\frac{y_B}{y_c}}}{y_B(1+y_B)^3} = \frac{f_{yy}}{z_B} < 0, \\ h_{yyy} &= -\frac{6(\alpha_2+\lambda_2)z_B}{(1+y_B)^4} = -\frac{3h_{yy}}{1+y_B} < 0, \\ h_{yyz} &= \frac{2(\alpha_2+\lambda_2)z_B}{(1+y_B)^3} = -\frac{3h_{yy}}{1+y_B} > 0. \end{aligned} \tag{38}$$

Thus, it follows that $F_{11}^1 < 0$, while both F_{12}^1 and F_{111}^1 are positive.

Since

$$\left(\frac{\delta}{4w^2+\delta^2} + \frac{1}{\delta}\right)(F_{11}^1)^2 + \left[\frac{4w^2}{\delta(4w^2+\delta^2)} + \frac{2\delta}{4w^2+\delta^2}\right](F_{12}^1)^2 \geq -\frac{4(2w^2+\delta^2)}{\delta(4w^2+\delta^2)}F_{11}^1F_{12}^1,$$

which together with $w^2 = \lambda_1\alpha_3$ produces that

$$\begin{aligned} &\left(\frac{\delta}{4w^2+\delta^2} + \frac{1}{\delta}\right)(F_{11}^1)^2 + \left[\frac{4w^2}{\delta(4w^2+\delta^2)} + \frac{2\delta}{4w^2+\delta^2}\right](F_{12}^1)^2 + \frac{w(8w^2+3\delta^2)}{\delta^2(4w^2+\delta^2)}F_{11}^1F_{12}^1 \\ &\geq -\frac{4(2w^2+\delta^2)}{\delta(4w^2+\delta^2)} - \frac{w(8w^2+3\delta^2)}{\delta^2(4w^2+\delta^2)}F_{11}^1F_{12}^1 = -\frac{4\delta(2w^2+\delta^2)-w(8w^2+3\delta^2)}{\delta^2(4w^2+\delta^2)}F_{11}^1F_{12}^1 \\ &\geq -\frac{8w(2w^2+\delta^2)-w(8w^2+3\delta^2)}{\delta^2(4w^2+\delta^2)}F_{11}^1F_{12}^1 > 0, \end{aligned} \tag{39}$$

where $\delta = \lambda_1 + \alpha_3$.

The sign of F_{11}^2 is determined by

$$\begin{aligned} \text{sign}\{F_{11}^2\} &= \text{sign}\{\lambda_1^2[\lambda_1(\alpha_3 - a)b^{-1}f_{yy} + (\alpha_3 + \lambda_1)h_{yy}]\} \\ &= \text{sign}\left\{-\frac{\lambda_1(\alpha_3 - a)(1 + 2y_B)}{y_B^2(1 + y_B)^2} + \frac{2(\alpha_2 + \lambda_2)(\alpha_3 + \lambda_1)}{(1 + y_B)^3}\right\} \\ &= \text{sign}\{2(\alpha_2 + \lambda_2)(\alpha_3 + \lambda_1)y_B^2 - \lambda_1(\alpha_3 - a)(1 + y_B)(1 + 2y_B)\} \\ &= \text{sign}\{2(\alpha_3 + \lambda_1)\lambda_2^2 - \lambda_1(\alpha_3 - a)(\alpha_2 + 2\lambda_2)\} \\ &= \text{sign}\{[2\lambda_2^2 - \lambda_1(\alpha_2 + 2\lambda_2)]\alpha_3 + 2\lambda_1\lambda_2^2 + a\lambda_1(\alpha_2 + 2\lambda_2)\}. \end{aligned} \tag{40}$$

Since $0 < \lambda_1 < \alpha_2y_B/(1 + y_B) = \alpha_2\lambda_2/(\alpha_2 + \lambda_2)$, we get that $2\lambda_2^2 - \lambda_1(\alpha_2 + 2\lambda_2) > 2\lambda_2^2 - \frac{\alpha_2\lambda_2}{\alpha_2 + \lambda_2}(\alpha_2 + 2\lambda_2) = \frac{\lambda_2}{\alpha_2 + \lambda_2}(2\lambda_2^2 - \alpha_2^2)$, which is nonnegative when the hypothesis

$\alpha_2^2 \leq 2\lambda_2^2$ holds. If $y_c = y_B(1 + y_B)$, then $a = 0$ from (32), and the sign of the last equation in (40) is positive. As a consequence, $F_{11}^2 > 0$.

Since $F_{12}^1 = -c\lambda_1^2 \Delta^{-1} h_{yz} = \frac{c\lambda_1^2(\alpha_2 + \lambda_2)}{\Delta(1 + y_B)^2}$, using (32) and the Hopf bifurcation condition (33), we have

$$\begin{aligned} & \text{sign}\left\{\frac{\delta(6w^2 + \delta^2)}{w^2(4w^2 + \delta^2)}F_{12}^1F_{11}^2 - \frac{c\lambda_1}{wz_B}F_{11}^2\right\} \\ = & \text{sign}\left\{\frac{\delta(6w^2 + \delta^2)}{w^2(4w^2 + \delta^2)}\frac{\lambda_1(\alpha_2 + \lambda_2)}{(1 + y_B)^2} - \frac{\Delta}{wz_B}\right\} = \text{sign}\left\{\frac{\delta(6w^2 + \delta^2)}{w^2(4w^2 + \delta^2)}\frac{\lambda_1(\alpha_2 + \lambda_2)}{(1 + y_B)^2} - \frac{w^2 + \delta^2}{bz_B}\right\} \\ = & \text{sign}\left\{\frac{\delta(6w^2 + \delta^2)}{w^2(4w^2 + \delta^2)}\frac{\lambda_1(\alpha_2 + \lambda_2)}{(1 + y_B)^2} - \frac{(w^2 + \delta^2)\alpha_2}{\delta(\alpha_3 - a)(1 + y_B)}\right\} = \text{sign}\left\{\frac{\delta(6w^2 + \delta^2)\lambda_1}{w^2(4w^2 + \delta^2)} - \frac{w^2 + \delta^2}{\delta(\alpha_3 - a)}\right\} \\ = & \text{sign}\left\{\left(\frac{\delta}{w^2} + \frac{2\delta}{4w^2 + \delta^2}\right)\lambda_1 - \frac{w^2 + \delta^2}{\delta(\alpha_3 - a)}\right\} = \text{sign}\left\{-\frac{\lambda_1}{\delta} + \frac{2\delta\lambda_1}{4w^2 + \delta^2}\right\} \\ = & \text{sign}\{\delta^2 - 4w^2\}. \end{aligned} \tag{41}$$

It is easy to see that $\delta^2 - 4w^2 = (\lambda_1 + \alpha_3)^2 - 4\lambda_1(\alpha_3 - a) = (\lambda_1 - \alpha_3)^2 \geq 0$. Therefore, (41) has a nonnegative sign.

The remaining terms which need to be proved in (37) include $\frac{2\delta}{4w^2 + \delta^2}(F_{12}^1)^2 + \frac{1}{\delta}(F_{11}^1)^2 - \frac{6w^2 + \delta^2}{w(4w^2 + \delta^2)}F_{11}^1F_{11}^2 - F_{111}^1$. Owing to

$$\begin{aligned} & \Delta^2\left[\frac{1}{\delta}(F_{11}^1)^2 + \frac{2\delta}{4w^2 + \delta^2}(F_{12}^1)^2 - \frac{6w^2 + \delta^2}{w(4w^2 + \delta^2)}F_{11}^1F_{11}^2\right] \\ = & \frac{1}{\delta}[\lambda_1^2(c\lambda_1w^{-1}f_{yy} - wh_{yy})]^2 + \frac{2\delta}{4w^2 + \delta^2}[-c\lambda_1^2h_{yz}]^2 \\ & - \frac{6w^2 + \delta^2}{w(4w^2 + \delta^2)}[\lambda_1^2(c\lambda_1w^{-1}f_{yy} - wh_{yy})]\lambda_1^2[\lambda_1(\alpha_3 - a)b^{-1}f_{yy} + (\alpha_3 + \lambda_1)h_{yy}] \\ = & \frac{\lambda_1^4z_B^2}{\delta w^2y_B^4(1 + y_B)^6}[\lambda_1^2\alpha_2(1 + 2y_B)x_B + 2w^2(\alpha_2 + \lambda_2)y_B^2]^2 + \frac{2\delta\lambda_1^4\alpha_2^2z_B^2(\alpha_2 + \lambda_2)^2}{(4w^2 + \delta^2)(1 + y_B)^6} \\ & + \frac{(6w^2 + \delta^2)\lambda_1^4z_B^2}{w^2(4w^2 + \delta^2)y_B^4(1 + y_B)^6}[\lambda_1^2\alpha_2(1 + 2y_B)x_B + 2w^2(\alpha_2 + \lambda_2)y_B^2][2\delta(\alpha_2 + \lambda_2)y_B^2 \\ & - w^2(1 + y_B)(1 + 2y_B)] \\ = & \frac{w^2\lambda_1^4z_B^2}{\delta y_B^4(1 + y_B)^6}[\delta(1 + y_B)(1 + 2y_B) + 2(\alpha_2 + \lambda_2)y_B^2]^2 + \frac{2\delta\lambda_1^4\alpha_2^2z_B^2(\alpha_2 + \lambda_2)^2}{(4w^2 + \delta^2)(1 + y_B)^6} \\ & + \frac{(6w^2 + \delta^2)\lambda_1^4z_B^2}{(4w^2 + \delta^2)y_B^4(1 + y_B)^6}[\delta(1 + y_B)(1 + 2y_B) + 2(\alpha_2 + \lambda_2)y_B^2][2\delta(\alpha_2 + \lambda_2)y_B^2 \\ & - w^2(1 + y_B)(1 + 2y_B)] \\ = & \frac{\lambda_1^4z_B^2}{y_B^4(1 + y_B)^6}\left\{\frac{w^2}{\delta}[\delta(1 + y_B)(1 + 2y_B) + 2(\alpha_2 + \lambda_2)y_B^2]^2 + \frac{2\delta\alpha_2^2(\alpha_2 + \lambda_2)^2y_B^4}{4w^2 + \delta^2}\right. \\ & \left. + \frac{6w^2 + \delta^2}{4w^2 + \delta^2}[\delta(1 + y_B)(1 + 2y_B) + 2(\alpha_2 + \lambda_2)y_B^2][2\delta(\alpha_2 + \lambda_2)y_B^2\right. \\ & \left. - w^2(1 + y_B)(1 + 2y_B)]\right\} \end{aligned} \tag{42}$$

and

$$\begin{aligned} \Delta^2F_{111}^1 &= \Delta\lambda_1^3(c\lambda_1w^{-1}f_{yyy} - wh_{yyy}) = \frac{\lambda_1^3(w^2 + \delta^2)}{b}(c\lambda_1f_{yyy} - w^2h_{yyy}) \\ &= \frac{\lambda_1^4\alpha_2(w^2 + \delta^2)z_B^2}{\delta y_B^3(1 + y_B)^5}[\delta(1 + y_B)(2 + 6y_B + 6y_B^2) + 6(\alpha_2 + \lambda_2)y_B^3] \\ &= \frac{\lambda_1^4z_B^2(w^2 + \delta^2)(\alpha_2 + \lambda_2)y_B}{\delta y_B^4(1 + y_B)^6}[\delta(1 + y_B)(2 + 6y_B + 6y_B^2) + 6(\alpha_2 + \lambda_2)y_B^3], \end{aligned} \tag{43}$$

we obtain

$$\begin{aligned}
 & \text{sign}\left\{\frac{1}{\delta}(F_{11}^1)^2 + \frac{2\delta}{4w^2+\delta^2}(F_{12}^1)^2 - \frac{6w^2+\delta^2}{w(4w^2+\delta^2)}F_{11}^1F_{11}^2 - F_{111}^1\right\} \\
 = & \text{sign}\left\{\frac{w^2}{\delta}[\delta(1+y_B)(1+2y_B) + 2(\alpha_2 + \lambda_2)y_B^2]^2 + \frac{2\delta\alpha_2^2(\alpha_2+\lambda_2)^2}{4w^2+\delta^2}y_B^4 \right. \\
 & + \frac{6w^2+\delta^2}{4w^2+\delta^2}[\delta(1+y_B)(1+2y_B) \\
 & + 2(\alpha_2 + \lambda_2)y_B^2][2\delta(\alpha_2 + \lambda_2)y_B^2 - w^2(1+y_B)(1+2y_B)] \\
 & \left. - \frac{(w^2+\delta^2)(\alpha_2+\lambda_2)y_B}{\delta}[\delta(1+y_B)(2+6y_B+6y_B^2) + 6(\alpha_2 + \lambda_2)y_B^3]\right\} \\
 = & \text{sign}\left\{\frac{(\alpha_2+\lambda_2)^2y_B^4(\delta^2\alpha_2^2-\delta^4-w^2\delta^2-4w^4)}{\delta(4w^2+\delta^2)} + \frac{(\alpha_2+\lambda_2)(\delta^4+7w^2\delta^2+2w^4)(1+3y_B+2y_B^2)y_B^2}{4w^2+\delta^2} \right. \\
 & \left. - \frac{w^4\delta(1+y_B)^2(1+2y_B)^2}{4w^2+\delta^2} - (w^2 + \delta^2)(\alpha_2 + \lambda_2)(1+y_B)(1+3y_B+3y_B^2)y_B\right\} \\
 = & \text{sign}\left\{y_B^2[\lambda_2^2(\delta^2\alpha_2^2 - \delta^4 - w^2\delta^2 - 4w^4) + \delta(\alpha_2 + 2\lambda_2)(\delta^4 + 7w^2\delta^2 + 2w^4)] \right. \\
 & \left. - [\delta^2w^4(1+2y_B)^2 + \delta\lambda_2(\delta^4 + 5w^2\delta^2 + 4w^4)(1+3y_B+3y_B^2)]\right\} \\
 = & \text{sign}\{H_1 - H_2\}.
 \end{aligned}
 \tag{44}$$

According to the hypothesis, $H_1 - H_2 > 0$, therefore, uniting (39), (40) and (44), the proof of this lemma is completed. \square

Remark 4. The hypothesis $H_1 > H_2$ in Theorem 4.5 implies that y_B cannot increase uncontrollable, thus y_c is bounded, which indicates that the existence of a stable periodic orbit in (3) is satisfied on the precondition of the bound of lymphocyte cells at the second stage. If the number of lymphocyte cells is much more than that of the cancer cells, that is to say, the function of the immune system is strong enough, then it is impossible for the cancer cells to survive stably for a long time.

5. Simulations. In this section we choose some suitable parameters in (4) to numerically simulate the theoretical conclusions obtained in the previous sections by using the software XPP.

(1) Choose $\alpha_1 = 500$, $\alpha_2 = \lambda_2 = 2$, $\lambda_1 = 0.5 < \frac{\alpha_2\lambda_2}{\alpha_2+\lambda_2}$, $x_0 = 0.125$, then $\alpha_3 = 0.25$ can be obtained from (9). It is easy to calculate that $x_B = \alpha_3\lambda_2/(\alpha_2\lambda_1) - x_0 = 0.375$, $y_B = \lambda_2/\alpha_2 = 1$, $z_B = \lambda_1x_B(1+y_B)\exp(y_B/y_c)/(\alpha_1y_B) = 0.001236$, $y_c = y_B(1+y_B) = 2$, $x_0^* = 0.125 = x_0$, which corresponds to the second case in Figure 2. All eigenvalues of the degenerate positive equilibrium B after perturbation have negative real parts at $\mu \neq 0$, thus point B is locally asymptotically stable and no Hopf bifurcation occurs in (3). Certainly periodic oscillation phenomena will not occur in the development of the immune system and cancer cells in (3) in this case.

The numerical simulations in Figures 3-4 support our conclusions when $\mu = 0$. Figure 3 describes the solution curve initiating from point (4, 10, 1.5) in the xyz -space. Figure 4 describe the three coordinate components in terms of time t of the solution curve in Figure 3, where (a), (b) and (c) represent the curves of $x(t)$, $y(t)$, $z(t)$, respectively.

(2) Choose $\alpha_1 = 1$, $\alpha_2 = \lambda_2 = 2$, $\lambda_1 = 0.5 < \frac{\alpha_2\lambda_2}{\alpha_2+\lambda_2}$, $x_0 = 0.0625$, $\alpha_3 = 0.07322$ can be obtained from (9), therefore, $x_B = 0.08394$, $y_B = \lambda_2/\alpha_2 = 1$, $z_B = 0.1384$, $y_c = y_B(1+y_B) = 2$ and $x_0^* = 0.0231 \neq x_0$. In addition, $w^2 = \alpha_3\lambda_1 = 0.03661$, $\delta = \alpha_3 + \lambda_1 = 0.57322$, $H_1 = 4.86176$ and $H_2 = 1.39616$ evidently satisfy the hypotheses in Theorem 4.5. A stable limit cycle will bifurcate from point B by perturbing the value of parameter α_3 near 0.0732, which means that there is a stable periodic orbit in \mathbb{R}^3 in the 2-stage cancer model (3) by choosing suitable parameters. Hence, the immune system and cancer cells can coexist for a quite long time on some initial conditions and their development exhibits stable periodic oscillation phenomenon.




Figure 3. Trajectories tend to the equilibrium point B in \mathbb{R}^3 as t increases when $\mu = 0$ and $x_0 = x_0^*$.




Figure 4. The plots of the coordinate components (a) $x(t)$, (b) $y(t)$, and (c) $z(t)$ of the solution in terms of time t near the equilibrium B when $\mu = 0$.




Figure 5. Three different solution orbits in the xyz -space with $\mu = 0.0051$ and $x_0 \neq x_0^*$, here the initial values are chosen as (a) $(0.1, 1.3, 0.3)$ (outside the limit cycle), (b) $(0.194, 1.184, 0.27)$ (on the limit cycle), and (c) $(0.09, 1.1, 0.1)$ (inside the limit cycle).

Figure 5 shows three solution orbits near the equilibrium point B on different initial conditions in \mathbb{R}^3 as $\mu = 0.0051$, where initial values are chosen as (a) $(0.1, 1.3, 0.3)$ (outside the limit cycle), (b) $(0.194, 1.184, 0.27)$ (on the limit cycle), and (c) $(0.09, 1.1, 0.1)$ (inside the limit cycle), respectively. For the sake of convenience, we denote the limit cycle by L . It can be observed that the degenerate equilibrium point B becomes a stable equilibrium point in this case.




Figure 6. The projections of the periodic orbit L on the (a) xy -, (b) yz -, and (c) xz -planes, respectively when $\mu = 0.0051$ and $x_0 \neq x_0^*$.




Figure 7. The plots of the components (a) $x(t)$, (b) $y(t)$, and (c) $z(t)$, respectively, of the periodic orbit in terms of the time t when $\mu = 0.0051$ and $x_0 \neq x_0^*$.

Figure 6 presents the projections of the periodic solution curve on the coordinate planes, where (a), (b) and (c) represent the projections on the xy -plane, yz -plane, xz -plane, respectively. Figure 7 corresponds to diagrams between the coordinate components and time t of the solution curve in Figure 6, where (a), (b) and (c) represent the curves of $x(t)$, $y(t)$, $z(t)$, respectively.

6. Discussion. Periodic oscillations with various periods have been observed in some cancer data (Fortin and Mackey [10]) and oscillations are common in the immune system (Stark et al. [29]). So it is reasonable to expect that the oscillatory phenomenon occurs in tumor and immune system interaction models. Rescigno and DeLisi [26] proposed a three equation model (3) to describe two different stages of lymphocytes in the interactions of the solid tumor and immune system. The oscillations existing in the signs of the derivatives of L_1 , L_2 and C subject to the time t were observed.

In this paper we established the existence and stability of periodic oscillations in the two-stage cancer model (3) by choosing the parameters suitably. After making a series of transformations of the variables, the qualitative analysis and some bifurcation results near the degenerate equilibrium were given. We have shown that the system could undergo Hopf bifurcation in the small neighborhood of the interior degenerate equilibrium. It is valuable to find out that any degenerate Hopf bifurcations cannot occur in this model, which excludes more complex dynamical behaviors.

The conditions on the stability of the bifurcated periodic orbits guarantee the appearance of stable periodic oscillations of the tumor levels. When a stable periodic orbit exists, it can be seen as “safe” in the neighborhood of this closed orbit since the trajectories originating from there will overwhelmingly go towards it. Therefore,




Figure 8. Trajectories spiral away from the unstable equilibrium point B when $a_1a_2 - a_3 < 0$.




Figure 9. The components (a) $x(t)$, (b) $y(t)$, and (c) $z(t)$ of the orbit in Figure 8 in terms of the time t when $a_1a_2 - a_3 < 0$.

the solid tumor and the immune system can coexist for a long term although the cancer is not eliminated. The oscillatory dynamics in the tumor and immune system interaction models demonstrate the phenomenon of long-term tumor relapse and have been observed in some related tumor and immune system models (d’Onofrio [7], Eftimie et al. [9], Kirschner and Panetta [15], Kuznetsov et al. [17], and Lejeune et al. [19]). We can interpret this situation from the biological point of view that while the immune system fights with cancer in the host, there exists a balance between them because of the periodic changes in the internal tissues and the external circumstances such that they coexist in a bounded region.

It is necessary to mention at the end that our discussions on the equilibrium point B are based on the hypothesis (9). If we choose parameters in (4) such that the left hand side term $a_1a_2 - a_3$ in (9) is positive, then B turns to be an asymptotically stable equilibrium point, which is similar to the first case of the numerical simulations in the last section. If $a_1a_2 - a_3 < 0$, then we find that B is an unstable nondegenerate equilibrium by numerical simulations (see Figures 8-9). In this case, the cancer cells will increase uncontrollably because their development are not controlled by the immune system any longer.

It will be very interesting to include immunotherapy in the tumor and immune system interaction model (Smyth et al. [28]), study the effects of adoptive cellular immunotherapy on the model, and explore conditions under which the tumor can

be eliminated (de Pillis et al. [6], Kirschner and Panetta [15]). We leave this for future consideration.

Finally, we would like to mention that in model (3) the development from the immature stage to the mature stage of lymphocytes was modeled by two ordinary differential equations in $L_1(t)$ and $L_2(t)$ explicitly. As pointed out by the reviewer, this process can be described by a time delay $\tau > 0$. Thus, we would like to propose the following two-stage cancer model described by two differential equations with a time delay:

$$\begin{aligned}\frac{dL}{dt} &= \lambda_1 L(t - \tau) - \alpha_3 L(t) + \alpha_1 \frac{L(t)C(t)}{1+KL(t)} \exp\left(-\frac{L(t)}{L_c}\right), \\ \frac{dC}{dt} &= \frac{\lambda_2 - \alpha_2 L(t)}{1+KL(t)} C(t),\end{aligned}\tag{45}$$

where the time delay $\tau > 0$ is the time for immature lymphocytes to develop into mature ones and $L(t)$ represents the density of the mature lymphocytes at time t . Recently, delayed models of tumor and immune response interactions have been studied extensively, we refer to d’Onofrio et al. [8], Bi and Ruan [2] and the references cited therein. In particular, Bi and Ruan [2] have shown that various bifurcations, including Hopf bifurcation, Bautin bifurcation and Hopf-Hopf bifurcation, can occur in such models. It would be interesting to consider the nonlinear dynamics of the delayed model (45).

REFERENCES

- [1] A. Albert, M. Freedman and A. S. Perelson, *Tumor and the immune system: The effects of tumor growth modulator*, Math. Biosci., **50** (1980), 25–58.
- [2] P. Bi and S. Ruan, *Bifurcations in tumor and immune system interaction models with delay effect*, submitted.
- [3] P. Boyle, A. d’Onofrio, P. Maisonneuve, G. Severi, C. Robertson, M. Tubiana and U. Veronesi, *Measuring progress against cancer in Europe: has the 15% decline targeted for 2000 come about?* Ann. Oncol., **14** (2003), 1312–1325.
- [4] E. A. Coddington and N. Levinson, “Theory of Ordinary Differential Equations,” McGraw-Hill Book Company, Inc., New York-Toronto-London, 1955.
- [5] C. DeLisi and A. Rescigno, *Immune surveillance and neoplasia-I: A minimal mathematical model*, Bull. Math. Biol., **39** (1977), 201–221.
- [6] L. G. de Pillis, A. E. Radunskaya, and C. L. Wiseman, *A validated mathematical model of cell-mediated immune response to tumor growth*, Cancer Res., **65** (2005), 7950–7958.
- [7] A. d’Onofrio, *A general framework for modeling tumor-immune system competition and immunotherapy: Mathematical analysis and biomedical inferences*, Physica D, **208** (2005), 220–235.
- [8] A. d’Onofrio, F. Gatti, P. Cerrai and L. Freschi, *Delay-induced oscillatory dynamics of tumour immune system interaction*, Math. Comput. Modelling, **51** (2010), 572–591.
- [9] R. Eftimie, J. L. Bramson and D.J.D. Earn, *Interactions between the immune system and cancer: A brief review of non-spatial mathematical models*, Bull. Math. Biol., **73** (2010), 2–32.
- [10] P. Fortin and M. C. Mackey, *Periodic chronicmyelogenous leukemia: Spectral analysis of blood cell counts and etiological implications*, Brit. J. Haematol., **104** (1999), 336–345.
- [11] K. O. Friedrichs, “Advanced Ordinary Differential Equations”, Gordon and Breach Science Publishers, New York, 1965.
- [12] M. Golubitsky and W. F. Langford, *Classification and unfoldings of degenerate Hopf bifurcations*, J. Differential Equations, **41** (1981), 375–415.
- [13] E. Hopf, *Abzweigung einer periodischen Lösung von einer stationären Lösung eines Differentialsystems*, Ber. Verh. Saechs. Akad. Wiss. Leipzig, Math. Naturwiss., **94** (1942), 3–22.
- [14] I.-D. Hsü and N. D. Kazarinoff, *An applicable Hopf bifurcation formula and instability of small periodic solutions of the Field-Noyes model*, J. Math. Anal. Appl., **55** (1976), 61–89.
- [15] D. Kirschner and J. Panetta, *Modeling immunotherapy of the tumor-immune interaction*, J. Math. Biol., **37** (1998), 235–252.

- [16] D. Kirschner and A. Tsygvintsev, *On the global dynamics of a model for tumor immunotherapy*, Math. Biosci. Engin., **6** (2009), 573–583.
- [17] V. A. Kuznetsov, I. A. Makalkin, M. A. Talor, and A. S. Perelson, *Nonlinear dynamics of immunogenic tumors: Parameters estimation and global bifurcation analysis*, Bull. Math. Biol., **56** (1994), 295–321.
- [18] R. Lefever and R. Garay, *Local discription of immune tumor rejection*, in “*Biomathematics and Cell Kinetics*” (eds. A. J. Valleron and P. D. M. Macdonald), North-Holland Biomedical Press, 1978, pp. 333–344.
- [19] O. Lejeune, M. A. J. Chaplain and I. El Akili, *Oscillations and bistability in the dynamics of cytotoxic reactions mediated by the response of immune cells to solid tumours*, Math. Computer Model., **47** (2008), 649–662.
- [20] A.-H. Lin, *A model of tumor and lymphocyte interactions*, Discrete. Contin. Dynam. Systems-B, **4** (2004), 241–266.
- [21] D. Liu, S. Ruan and D. Zhu, *Bifurcation analysis in models of tumor and immune system interactions*, Discrete Contin. Dynam. Systems-B, **12** (2009), 151–168.
- [22] W. Liu, T. Hillen and H. I. Freedman, *A mathematical model for M-phase specific chemotherapy including the G_0 -phase and immunoresponse*, Math. Biosci. Engin., **4** (2007), 239–259.
- [23] Z. Lu and Y. Luo, *Two limit cycles in three-dimensional Lotka-Volterra systems*, Comput. Math. Appl., **44** (2002), 51–66.
- [24] L. Perko, “*Differential Equations and Dynamical Systems*”, 3rd edition, Texts in Applied Mathematics, 7, Springer, New York, 2001.
- [25] A. B. Poore, *On the theory and application of the Hopf-Friedrichs bifurcation theory*, Arch. Rational Mech. Anal., **60** (1976), 371–393.
- [26] A. Rescigno and C. DeLisi, *Immune surveillance and neoplasia-II: A two-stage mathematical model*, Bull. Math. Biol., **39** (1977), 487–497.
- [27] H. L. Smith, “*Monotone Dynamical Systems: An Introduction to Theory of Competitive and Cooperative Systems*”, Math. Surveys Monogr., **41**, Amer. Math. Soc., Providence, RI, 1995.
- [28] M. J. Smyth, D. I. Godfrey and J. A. Trapani, *A fresh look at tumor immunosurveillance and immunotherapy*, Nat. Immunol., **4** (2001), 293–299.
- [29] J. Stark, C. Chan and A. J. T. George, *Oscillations in the immune system*, Immunol. Rev., **216** (2007), 213–231.
- [30] G. W. Swan, *Immunological surveillance and neoplastic development*, Rocky Mountain J. Math., **9** (1979), 143–148.
- [31] R. Thomlinson, *Measurement and management of carcinoma of the breast*, Clin. Radiol., **33** (1982), 481–492.
- [32] J. Zhang, “*The Geometric Theory and Bifurcation Problem of Ordinary Differential Equation*”, Peking University Press, Beijing, 1987.

Received January 28, 2011; Accepted October 6, 2011.

E-mail address: liudan@xidian.edu.cn

E-mail address: ruan@math.miami.edu

E-mail address: dmzhu@math.ecnu.edu.cn