

1. Suppose that a fair coin is flipped 6 times in sequence and let X be the number of “heads” that show up. Draw Pascal’s triangle down to the sixth row (recall that the zeroth row consists of a single 1) and use your table to compute the probabilities $P(X = k)$ for $k = 0, 1, 2, 3, 4, 5, 6$.

Here is Pascal’s Triangle:

$$\begin{array}{cccccccc}
 & & & & & & & 1 \\
 & & & & & & 1 & 1 \\
 & & & & 1 & 2 & 1 & \\
 & & 1 & 3 & 3 & 1 & & \\
 & 1 & 4 & 6 & 4 & 1 & & \\
 1 & 5 & 10 & 10 & 5 & 1 & & \\
 1 & 6 & 15 & 20 & 15 & 6 & 1 &
 \end{array}$$

Then since $2^6 = 64$ we have the following table of probabilities:

k	0	1	2	3	4	5	6
$P(X = k)$	$\frac{1}{64}$	$\frac{6}{64}$	$\frac{15}{64}$	$\frac{20}{64}$	$\frac{15}{64}$	$\frac{6}{64}$	$\frac{1}{64}$

2. Suppose that a fair coin is flipped 4 times in sequence.

- List all 16 outcomes in the sample space S .
- List the outcomes in each of the following events:
 $A = \{\text{at least 3 heads}\}$,
 $B = \{\text{at most 2 heads}\}$,
 $C = \{\text{heads on the 2nd flip}\}$,
 $D = \{\text{exactly 2 tails}\}$.
- Assuming that all outcomes are **equally likely**, use the formula $P(E) = \#E/\#S$ to compute the following probabilities:

$$P(A \cup B), \quad P(A \cap B), \quad P(C), \quad P(D), \quad P(C \cap D).$$

(a) The sample space is

$$\begin{aligned}
 S = \{ & HHHH, \\
 & HHHT, HHTH, HTHH, THHH, \\
 & HHTT, HTHT, HTTH, THHT, THTH, TTHH, \\
 & HTTT, THTT, TTHT, TTTH, \\
 & TTTT \}
 \end{aligned}$$

(b) The events are

$$\begin{aligned}
 A &= \{HHHH, \\
 &\quad HHHT, HHTH, HTTH, THHH\}, \\
 B &= \{HHTT, HTHT, HTTH, THHT, THTH, TTTH, \\
 &\quad HTTT, THTT, TTHT, TTTT\}, \\
 C &= \{HHHH, \\
 &\quad HHHT, HHTH, THHH, \\
 &\quad HHTT, THHT, THTH, \\
 &\quad THTT\}, \\
 D &= \{HHTT, HTHT, HTTH, THHT, THTH, TTTH\}.
 \end{aligned}$$

(c) Observe that $A \cup B = S$ and $A \cap B = \emptyset$, so that

$$P(A \cup B) = P(S) = 1 \quad \text{and} \quad P(A \cap B) = P(\emptyset) = 0.$$

Observe that $\#C = 8$ and $\#D = 6$, so that

$$P(C) = \frac{\#C}{\#S} = \frac{8}{16} \quad \text{and} \quad P(D) = \frac{\#D}{\#S} = \frac{6}{16}.$$


Finally, observe that $C \cap D = \{HHTT, THHT, THTH\}$ so that

$$P(C \cap D) = \frac{\#(C \cap D)}{\#S} = \frac{3}{16}.$$

3. Draw Venn diagrams to verify *de Morgan's laws*: For all events $E, F \subseteq S$ we have

- (a) $(E \cup F)' = E' \cap F'$,
- (b) $(E \cap F)' = E' \cup F'$.

The proof follows from the following diagrams:



4. Suppose that a fair coin is flipped until heads appears. The sample space is

$$S = \{H, TH, TTH, TTTH, TTTTH, \dots\}.$$

However these outcomes are **not equally likely**.

- (a) Let E_k be the event {first H occurs on the k th flip}. Explain why $P(E_k) = 1/2^k$.
 [Hint: The outcomes of the coin flips are **independent**.]
- (b) Use the “geometric series” to verify that the sum of all the probabilities equals 1:

$$\sum_{k=1}^{\infty} P(E_k) = 1.$$

(a) There is exactly one outcome in this event:

$$E_k = \{\underbrace{TTT \cdots T}_{k-1 \text{ times}} H\}.$$

Since the coin flips are fair and independent we have

$$\begin{aligned} P(E_k) &= P(\underbrace{TTT \cdots T}_{k-1 \text{ times}} H) \\ &= \underbrace{P(T)P(T)P(T) \cdots P(T)}_{k-1 \text{ times}} P(H) \\ &= \underbrace{\left(\frac{1}{2}\right) \left(\frac{1}{2}\right) \left(\frac{1}{2}\right) \cdots \left(\frac{1}{2}\right)}_{k-1 \text{ times}} \left(\frac{1}{2}\right) = \left(\frac{1}{2}\right)^k = \frac{1}{2^k}. \end{aligned}$$

(b) Recall the “geometric series” from calculus: If q is any number satisfying $-1 < q < 1$ then we have

$$1 + q + q^2 + q^3 + \cdots = \frac{1}{1-q}.$$

By substituting $q = 1/2$ we obtain

$$\begin{aligned} 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \cdots &= \frac{1}{1 - 1/2} \\ 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \cdots &= 2 \\ \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \cdots &= 2 - 1 \\ \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \cdots &= 1 \end{aligned}$$

and hence

$$\sum_{k=0}^{\infty} P(E_k) = \sum_{k=0}^{\infty} \frac{1}{2^k} = \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \cdots = 1.$$

5. Suppose that $P(A) = 0.5$, $P(B) = 0.6$ and $P(A \cap B) = 0.3$. Use this information to compute the following probabilities. A Venn diagram may be helpful.

- (a) $P(A \cup B)$,
- (b) $P(A \cap B')$,
- (c) $P(A' \cup B')$.

(a) Using Inclusion-Exclusion for two events gives

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) = 0.5 + 0.6 - 0.3 = \boxed{0.8}.$$

(b) Using the Law of Total Probability gives

$$\begin{aligned} P(A) &= P(A \cap B) + P(A \cap B') \\ 0.5 &= 0.3 + P(A \cap B') \\ \boxed{0.2} &= P(A \cap B'). \end{aligned}$$

(c) Using de Morgan's Law and Complementary Events gives

$$P(A' \cup B') = P((A \cap B)') = 1 - P(A \cap B) = 1 - 0.3 = \boxed{0.7}.$$

6. Let X be a real number that is selected randomly from $[0, 1]$, i.e., the closed interval from zero to one. Use your intuition to assign values to the following probabilities:

- (a) $P(X = 1/2)$,
- (b) $P(0 \leq X \leq 1/2)$,
- (c) $P(0 < X < 1/2)$,
- (d) $P(1/3 < X \leq 3/4)$,
- (e) $P(-1 < X < 3/4)$.

(a) If all of the points in $[0, 1]$ are "equally likely," then since there are infinitely many points we must have

$$P(X = 1/2) = \frac{1}{\infty} = 0.$$

Maybe you're uncomfortable with this, but it's the least wrong answer we can come up with.

(c) By symmetry, there must be a 50% of landing in the left half of the interval:

$$P(0 < X < 1/2) = 1/2.$$

(b) If you agreed in part (a) that $P(X = 1/2) = P(X = 0) = 0$ then we must have

$$\begin{aligned} P(0 \leq X \leq 1/2) &= \cancel{P(X=0)} + P(0 < X < 1/2) + \cancel{P(X=1/2)} \\ &= 0 + P(0 < X < 1/2) + 0 \\ &= P(0 < X < 1/2) = 1/2. \end{aligned}$$

(c) In general, the probability of landing in an interval must be the **length** of the interval. And we can just ignore the endpoints.

$$P(1/3 < X \leq 3/4) = \frac{3}{4} - \frac{1}{3} = \frac{5}{12}.$$

(d) It is impossible to get $-1 < X < 0$, so we must have

$$\begin{aligned} P(-1 < X < 3/4) &= \cancel{P(-1 < X < 0)} + P(0 < X < 3/4) \\ &= 0 + P(0 < X < 3/4) \\ &= P(0 < X < 3/4) = 3/4. \end{aligned}$$

7. Consider a strange coin with $P(H) = p$ and $P(T) = q = 1 - p$. Suppose that you flip the coin n times and let X be the number of heads that you get. Find a formula for the probability $P(X \geq 1)$. [Hint: Observe that $P(X \geq 1) + P(X = 0) = 1$. Maybe it's easier to find a formula for $P(X = 0)$.]

There is only one way to get $X = 0$:

$$“X = 0” = \underbrace{\{TTT \cdots T\}}_{n \text{ times}}.$$

Then by independence we must have

$$\begin{aligned} P(X = 0) &= P(\underbrace{TTT \cdots T}_{n \text{ times}}) \\ &= \underbrace{P(T)P(T)P(T) \cdots P(T)}_{n \text{ times}} \\ &= \underbrace{qqq \cdots q}_{n \text{ times}} = q^n \end{aligned}$$

and hence $P(X \geq 1) = 1 - P(X = 0) = 1 - q^n$.

8. Suppose that you roll a pair of fair six-sided dice.

- (a) Write down all elements of the sample space S . What is $\#S$? Are the outcomes equally likely? [Hopefully, yes.]
- (b) Compute the probability of getting a “double six.” [Hint: Let $E \subseteq S$ be the subset of outcomes that correspond to getting a “double six.” Assuming that the outcomes of your sample space are equally likely, you can use the formula $P(E) = \#E/\#S$.]

(a) Let's suppose that one die is “blue” and the other is “red,” so we can tell them apart. In other words, the outcome “12” = “the blue die shows 1 and the red die shows 2” will differ from

the outcome “21”=“the blue die shows 2 and the red die shows 1.” The the sample space is:

$$S = \{11, 12, 13, 14, 15, 16 \\ 21, 22, 23, 24, 25, 26 \\ 31, 32, 33, 34, 35, 36 \\ 41, 42, 43, 44, 45, 46 \\ 61, 62, 63, 64, 65, 66\}.$$

Independence and fairness suggest that for any outcome $ij \in S$ we must have $P(ij) = P(i)P(j) = (1/6)(1/6) = 1/36$. In other words, the 36 outcomes are equally likely.¹

(b) Let E = “double six,” so that $E = \{66\}$. Then we have

$$P(E) = \frac{\#E}{\#S} = \frac{1}{36}.$$

9. Analyze the Chevalier de Méré’s two experiments:

- (a) Roll a fair six-sided die 4 times and let X be the number of “sixes” that you get. Compute $P(X \geq 1)$. [Hint: You can think of a die roll as a “strange coin flip,” where H = “six” and T = “not six.” Use Problem 7.]
- (b) Roll a pair of fair six-sided dice 24 times and let Y be the number of “double sixes” that you get. Compute $P(Y \geq 1)$. [Hint: You can think of rolling two dice as a “very strange coin flip,” where H = “double six” and T = “not double six.” Use Problems 7 and 8.]

(a) Roll a fair six-sided die and let H = “we get six,” so that $P(H) = p = 1/6$ and $P(T) = q = 5/6$. Then according to Problem 7 we have

$$P(X \geq 1) = 1 - q^4 = 1 - \left(\frac{5}{6}\right)^4 = \boxed{51.77\%}.$$

(b) Roll a pair of fair six-sided dice and let H = “we get double six.” Then from Problem 8 we know that $P(H) = p = 1/36$ and $P(T) = q = 35/36$ and from Problem 7 we find

$$P(Y \geq 1) = 1 - q^{24} = 1 - \left(\frac{35}{36}\right)^{24} = \boxed{49.14\%}.$$

10. Roll a fair six-sided die three times in sequence, and consider the events

$$E_1 = \{\text{you get 1 or 2 or 3 on the first roll}\}, \\ E_2 = \{\text{you get 1 or 3 or 5 on the second roll}\}, \\ E_3 = \{\text{you get 2 or 4 or 6 on the third roll}\}.$$

You can assume that $P(E_1) = P(E_2) = P(E_3) = 1/2$.

- (a) Explain why $P(E_1 \cap E_2) = P(E_1 \cap E_3) = P(E_2 \cap E_3) = 1/4$ and $P(E_1 \cap E_2 \cap E_3) = 1/8$.
 (b) Use this information to compute $P(E_1 \cup E_2 \cup E_3)$.

¹It’s perfectly okay to consider the two dice as “unordered” or “uncolored.” Then we will have $\#S = 21$. However, in this case the outcomes will **not** be equally likely, which makes the analysis much harder.

(a) Since the event E_i only cares about what happens on the i th roll, we will assume that these events are **independent**. Then for all $i < j$ we have

$$P(E_i \cap E_j) = P(E_i)P(E_j) = \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4}$$

and for all $i < j < k$ we have

$$P(E_i \cap E_j \cap E_k) = P(E_i)P(E_j)P(E_k) = \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{8}.$$

(b) Now we can use the Principle of Inclusion-Exclusion:

$$\begin{aligned} P(E_1 \cup E_2 \cup E_3) &= P(E_1) + P(E_2) + P(E_3) \\ &\quad - P(E_1 \cap E_2) - P(E_1 \cap E_3) - P(E_2 \cap E_3) \\ &\quad + P(E_1 \cap E_2 \cap E_3) \\ &= 1/2 + 1/2 + 1/2 \\ &\quad - 1/4 - 1/4 - 1/4 \\ &\quad + 1/8 \\ &= 3/2 - 3/4 + 1/8 = \boxed{7/8}. \end{aligned}$$

Alternate Solution. We can think of this experiment as a “very strange coin flip,” in which the definition of “heads” changes from flip to flip:

$$\begin{aligned} E_1 &= \{\text{you get heads on the first flip}\}, \\ E_2 &= \{\text{you get heads on the second flip}\}, \\ E_3 &= \{\text{you get heads on the third flip}\}. \end{aligned}$$

Since the probability of “heads” is always $1/2$ and since the events are independent, we can treat this just like three flips of a fair coin. Then using Problem 7 gives

$$\begin{aligned} P(E_1 \cup E_2 \cup E_3) &= P(\text{you get heads at least once}) \\ &= 1 - P(\text{tails})^3 \\ &= 1 - (1/2)^3 = 7/8. \end{aligned}$$

This simplification seems a bit dubious but it must be okay because we got the correct answer.